



Fisheries and Oceans
Canada

Pêches et Océans
Canada

Science

Sciences

C S A S

Canadian Science Advisory Secretariat

S C C S

Secrétariat canadien de consultation scientifique

Research Document 2012/109

Document de recherche 2012/109

Pacific Region

Région du Pacifique

**Updated stock assessment for
Bocaccio (*Sebastes paucispinis*) in
British Columbia waters for 2012**

**Mise à jour de l'évaluation des stocks de
bocaccio (*Sebastes paucispinis*) dans
les eaux de la Colombie-Britannique
en 2012**

R. D. Stanley¹, M. McAllister², and P. Starr³

¹ Marine Ecosystem and Aquaculture Division,
Fisheries and Oceans Canada, Pacific Biological Station,
3190 Hammond Bay Rd.,
Nanaimo, BC V9T 6N7

² Fisheries Centre
Aquatic Ecosystems Research Laboratory (AERL)
2200 Main Mall, University of British Columbia
Vancouver, BC V6T 1Z4

³ Canadian Groundfish
Research and Conservation Society
1406 Rose Ann Drive,
Nanaimo, BC V9T 4K8

This series documents the scientific basis for the evaluation of aquatic resources and ecosystems in Canada. As such, it addresses the issues of the day in the time frames required and the documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

La présente série documente les fondements scientifiques des évaluations des ressources et des écosystèmes aquatiques du Canada. Elle traite des problèmes courants selon les échéanciers dictés. Les documents qu'elle contient ne doivent pas être considérés comme des énoncés définitifs sur les sujets traités, mais plutôt comme des rapports d'étape sur les études en cours.

Research documents are produced in the official language in which they are provided to the Secretariat.

Les documents de recherche sont publiés dans la langue officielle utilisée dans le manuscrit envoyé au Secrétariat.

This document is available on the Internet at:

Ce document est disponible sur l'Internet à:

<http://www.dfo-mpo.gc.ca/csas-sccs>

ISSN 1499-3848 (Printed / Imprimé)

ISSN 1919-5044 (Online / En ligne)

© Her Majesty the Queen in Right of Canada, 2012

© Sa Majesté la Reine du Chef du Canada, 2012

Canada

TABLE OF CONTENTS

| | |
|---|------|
| LIST OF TABLES..... | IV |
| LIST OF APPENDIX TABLES | V |
| LIST OF FIGURES | V |
| LIST OF APPENDIX FIGURES | VII |
| ABSTRACT..... | VIII |
| RÉSUMÉ..... | IX |
| 1 INTRODUCTION..... | 1 |
| 1.1 CONTEXT | 1 |
| 1.2 OBJECTIVES..... | 1 |
| 2 STOCK STRUCTURE..... | 2 |
| 3 DISTRIBUTION | 3 |
| 4 LIFE HISTORY PARAMETERS | 3 |
| 4.1 SAMPLE SOURCES..... | 3 |
| 4.2 GROWTH, AGE-AT-MATURITY, AND NATURAL MORTALITY | 4 |
| 5 CURRENT MANAGEMENT AND MONITORING OF BOCACCIO CATCHES..... | 7 |
| 6 COMMERCIAL CATCH DATA | 8 |
| 6.1 TRAWL CATCH | 8 |
| 6.2 HOOK-AND-LINE CATCH | 9 |
| 6.3 RECREATIONAL CATCH..... | 10 |
| 7 ABUNDANCE TRENDS - COMMERCIAL CPUE..... | 13 |
| 8 ABUNDANCE TRENDS - SURVEYS..... | 14 |
| 9 TRAWL SURVEY CATCHABILITY | 18 |
| 10 BAYESIAN SURPLUS PRODUCTION MODEL..... | 19 |
| 11 REFERENCE CASE..... | 24 |
| 11.1 REFERENCE CASE INPUTS | 24 |
| 11.2 REFERENCE CASE RESULTS..... | 24 |
| 11.3 REFERENCE CASE MODEL PERFORMANCE AND UNCERTAINTY | 31 |
| 11.4 REFERENCE CASE RECONSTRUCTION OF HISTORICAL CATCHES..... | 35 |
| 12 SENSITIVITY TESTS | 37 |
| 12.1 SENSITIVITY RUNS | 37 |
| 12.2 SENSITIVITY RESULTS..... | 40 |
| 12.3 SENSITIVITY RUNS EVALUATED WITH BAYES FACTORS | 46 |
| 13 DECISION TABLES | 48 |
| 14 STATUS OF BOCACCIO IN U.S. WATERS | 57 |
| 15 SUMMARY | 58 |
| 16 RESPONSES TO 2009 RECOMMENDATIONS FOR FUTURE WORK..... | 59 |
| 17 RECOMMENDATIONS FOR FUTURE WORK..... | 60 |
| 18 ACKNOWLEDGEMENTS..... | 60 |
| 19 LITERATURE CITED | 61 |
| 20 APPENDIX A CATCH..... | 64 |

| | | |
|----|---|----|
| 21 | APPENDIX B ESTIMATION OF ABUNDANCE INDICES FROM THE IPHC SURVEY | 67 |
| 22 | APPENDIX C RELATIVE ABUNDANCE INDICES | 68 |
| 23 | APPENDIX D ADDITIONAL REFERENCE CASE RESULTS..... | 72 |

LIST OF TABLES

| | |
|--|----|
| Table 1. Updated growth parameters for Bocaccio. | 5 |
| Table 2. The seven fisheries modelled in this stock assessment, showing the years these fisheries were included. | 8 |
| Table 3. Estimates of Bocaccio catch from Minor Area 12. | 11 |
| Table 4. Estimates of recreational catch of Bocaccio 2000-2010. | 11 |
| Table 5. Fishery independent surveys used in this assessment. | 15 |
| Table 6. Observations by year for the abundance indices used in the assessment. | 15 |
| Table 7. Survey catch rates (pieces/survey), frequency of occurrence, and mean lengths. | 15 |
| Table 8. Posterior means and CVs for q-gross (qgfin) for the current and 2009 assessments. | 19 |
| Table 9. Prior pdfs of parameters K, q for the commercial CPUE data, P_0 , and r. | 21 |
| Table 10. Key parameter choices for the current reference case and significant changes from the 2009 reference case. | 26 |
| Table 11. Reference case 2012 stock assessment statistics. | 27 |
| Table 12. Posterior 10th, 50th and 90 th percentiles of stock biomass (t) 1935-2012 from the reference case run. | 28 |
| Table 13. Posterior 10th, 50th and 90 th percentiles of the ratio of stock biomass in each recent year to the stock biomass sixty years prior taken from Table 12. | 29 |
| Table 14. Comparison of key results from the 2009 and 2012 analyses. | 29 |
| Table 15. Values for the CVs applied for each of the abundance indices in the reference case and other model runs. | 34 |
| Table 16. Summary of sensitivity test runs. | 38 |
| Table 17. Records of Pacific Halibut catch (t) in BC waters from 1900-1937. | 40 |
| Table 18. Medians and 80% credibility intervals drawn from the posterior distributions for seven parameters taken from the Bocaccio assessment for the reference run and 18 sensitivity runs. | 43 |
| Table 19. Posterior medians for estimated catch values and replacement yield statistics for 2012 taken from the Bocaccio assessment for the reference run and a selection of the sensitivity runs. | 44 |
| Table 20. Bayes factors for alternative mode runs. | 47 |
| Table 21. Decision tables are provided in the following runs for the reference case and five sensitivity runs. | 48 |
| Table 22. Stock status indicators for Bocaccio after 5, 20, and 60 years for the reference case. | 49 |
| Table 23. Stock status indicators for Bocaccio after 5, 20, and 60 years for Case B.1 low prior r mean. | 51 |
| Table 24. Stock status indicators for Bocaccio after 5, 20 and 60 years for Case B.2, high prior r mean. | 52 |
| Table 25. Stock status indicators for Bocaccio after 5, 20, and 60 years for Case A.1 B_{msy}/B_0 set at 0.3. | 53 |
| Table 26. Stock status indicators for Bocaccio after 5, 20, and 60 years for Case A.2 B_{msy}/B_0 set at 0.4. | 54 |
| Table 27. Stock status indicators for Bocaccio after 5, 20, and 60 years for Case A.3 B_{msy}/B_0 set at 0.6. | 55 |
| Table 28. Summary decision table for the probability that stock biomass exceeds $0.4*B_{msy}$ within 60 years under each alternative constant TAC policy (t) and under each alternative hypothesized prior mean value for the parameter for the maximum intrinsic rate of increase r. | 56 |
| Table 30. Summary decision table for the probability that stock biomass exceeds $0.4*B_{msy}$ within 60 years under each alternative constant TAC policy (t) and under each alternative hypothesized value for t B_{msy}/K | 56 |

| | |
|---|----|
| Table 31. Summary decision table for the probability that stock biomass exceeds $0.4 \cdot B_{msy}$ within 60 years under each alternative constant TAC policy (t) and under each alternative hypothesized scenario for the level of historic trawl and non-halibut hook and line catch. | 56 |
| Table 32. Decision table showing the time to reach four reference points (RP): $0.4 \cdot B_{msy}$, $0.8 \cdot B_{msy}$, $0.5 \cdot B_{t-3Gen}$, $0.7 \cdot B_{t-3Gen}$ over a range of constant catch quota policies (t) for two levels of confidence for the reference case run (see text for a description of these reference points). | 57 |

LIST OF APPENDIX TABLES

| | |
|--|----|
| Appendix Table 1. Inputted catch values in the Reference case (1935-1975). | 64 |
| Appendix Table 2. Inputted catches in the Reference case (1976-2012). | 65 |
| Appendix Table 3. Time series of fishery effort in the halibut, salmon troll, and recreational fisheries | 66 |
| Appendix Table 4. Arithmetic and standardised commercial bottom trawl CPUE indices with upper and lower bounds of the standardised indices and the associated standard error for the 3C-5E model of non-zero catches of Bocaccio. | 68 |
| Appendix Table 5. Biomass estimates for Bocaccio from the QCSd Shrimp Trawl Survey for the survey years 1999 to 2011. | 68 |
| Appendix Table 6. Biomass estimates for Bocaccio from the WCVI shrimp trawl survey for the survey years 1975 to 2011. | 69 |
| Appendix Table 7. Biomass estimates for Bocaccio from the West Coast Haida Gwaii groundfish synoptic trawl survey for the years 2006 to 2010. | 70 |
| Appendix Table 8. Biomass estimates for Bocaccio from the Hecate Strait Groundfish synoptic trawl survey for the years 2005 to 2011. | 70 |
| Appendix Table 9. Biomass estimates for Bocaccio from the Queen Charlotte Sound Groundfish synoptic trawl survey for the years 2005 to 2011. | 70 |
| Appendix Table 10. Biomass estimates for Bocaccio from the West Coast Vancouver Island Groundfish synoptic trawl survey for the years 2006 to 2010. | 71 |
| Appendix Table 11. Biomass estimates for Bocaccio in the U.S. Triennial survey (Canadian waters only) with 95% confidence regions based on the bootstrap distribution of biomass. | 71 |
| Appendix Table 12. Estimates and 95% confidence limits of relative catch rate (pieces/skate) of Bocaccio in the IPHC BC longline survey. | 71 |
| Appendix Table 13. Posterior means, medians, standard deviations (SD), CVs and 95% probability intervals for q-gross (qgfin). | 72 |

LIST OF FIGURES

| | |
|---|----|
| Figure 1. Area of occupancy for Bocaccio based on survey and commercial observations from 2002-2011. | 3 |
| Figure 2. Bubble plot of proportions-at-age for female and male Bocaccio for 2001-2010, all sources combined. | 4 |
| Figure 3. Observed and estimated length-at-age for a. female and b. male Bocaccio. | 6 |
| Figure 4. Observed and estimated proportion mature-at-age for female Bocaccio. | 6 |
| Figure 5. Female Bocaccio proportion mature-at-age (line) compared with histogram of age frequency of females, for all samples combined. | 7 |
| Figure 6. Domestic (U.S. and Can.) and foreign (Soviet, Japanese, and Polish) trawl landings (1930 to 2011). | 9 |
| Figure 7. Time series of annual effort used for imputation of total catch in the salmon troll, halibut, and recreational fisheries. | 10 |

| | |
|---|----|
| Figure 8. Plot of the initial prior distribution for recreational catch of Bocaccio in BC in 2010 from a qualitative interpretation based on recreational catch data and interviews..... | 13 |
| Figure 9. Comparison of the lognormal and binomial standardised commercial trawl CPUE indices for Bocaccio. | 14 |
| Figure 10. Locations of Shrimp trawl, US NMFS Triennial bottom trawl, and IPHC HL longline surveys. | 16 |
| Figure 11. Locations of Groundfish Synoptic bottom trawl surveys..... | 16 |
| Figure 12. Plots of biomass estimates for Bocaccio from the a. West Coast Vancouver Island shrimp trawl, b. Queen Charlotte Sound shrimp trawl, c. U.S. Triennial survey, and d. IPHC longline surveys. | 17 |
| Figure 13. Plots of biomass estimates for Bocaccio from the a. West Coast Haida Gwaii; b. Hecate Strait, c. Queen Charlotte Sound and, d. West Coast Vancouver Island Groundfish Synoptic trawl surveys for 2003 to 2011. | 18 |
| Figure 14. Fitted normal prior density function of the maximum intrinsic rate of increase, r , for BC Bocaccio. | 21 |
| Figure 15. Estimated values for B_{msy}/B_0 under the Ricker and Beverton-Holt stock-recruit models as a function of the steep parameter in the a) Ricker and b) Beverton-Holt stock-recruit functions..... | 23 |
| Figure 16. A plot of the reference case Schaefer and three alternative production functions applied in evaluations of the sensitivity of results to different model settings. | 24 |
| Figure 17. Plots of median and 80% probability intervals and indices rescaled by their median a. 1935-2012; b. 1975-2012; c. 2000-2012. | 30 |
| Figure 18. Estimated stock status in 2012 and exploitation rate relative to reference points..... | 31 |
| Figure 19. Marginal posterior distributions for a. carrying capacity (K or B_0); b. maximum sustainable yield (msy); c. r ; d. replacement yield in 2012; e. stock biomass in 2012; f. the ratio of fishing mortality rate in 2012 to that at msy ; g. the ratio of stock biomass in 2012 to average unfished stock size; and, h. the ratio of total catch biomass in 2012 to replacement yield in 2012. | 33 |
| Figure 20. Annual deviates in surplus production with the median and 80% probability intervals shown. | 34 |
| Figure 21. Median and 80% probability intervals for catch of Bocaccio in the: a. halibut fishery, b. salmon troll fishery, and c. recreational fishery, d. all sectors combined including trawl and ZN HL. | 36 |
| Figure 22. Prior and posterior density functions for the catchability coefficients for the a. halibut, b. salmon troll, and c. recreational sectors..... | 37 |
| Figure 23. Plots of a. effective halibut effort in BC waters and b. posterior median catch of Bocaccio by halibut gear under different scenarios for constant percent changes in catchability of halibut gear for Bocaccio, (i.e., termed “technological creep” or “TC”)...... | 45 |
| Figure 24. Plots of the a. marginal posterior distributions for total catch and replacement yield; b. draws from the posterior distribution of values for replacement yield versus total catch from the reference case run; c. marginal posterior distributions for total catch and replacement yield and d. draws from the posterior distributions of values for replacement yield versus total catch from run A.1..... | 46 |
| Figure 25. Reference case plots of the ratio of a. median stock biomass to B_{msy} for different constant total catch policies and b. 10 th , 50 th (median), and 90 th percentiles. | 50 |
| Figure 26. Estimated spawning output time series 1892-2011 for the base case, with approximate 95% confidence intervals. | 58 |

LIST OF APPENDIX FIGURES

Appendix Figure 1. Marginal density functions for q-gross (qgfin) for the seven different surveys
when Bayesian updating and uncertainty factors are applied to the q-net factors 73

Correct citation for this publication:

Stanley, R.D., McAllister, M. and Starr, P. 2012. Updated stock assessment for Bocaccio (*Sebastes paucispinis*) in British Columbia waters for 2012. DFO Can. Sci. Adv. Sec. Res. Doc. 2012/109. ix + 73 p.

ABSTRACT

This document provides a stock assessment for Bocaccio in British Columbia waters using data current to 2011. Results of the work are intended to serve as advice over the short term to managers and stakeholders on current stock status, and likely impacts of different harvest options. As in previous work, a Bayesian surplus production model was used. It was fit to one fishery dependent and eight fishery independent biomass indices, and a reconstructed catch history back to 1935 when the population was assumed to be near to an unfished equilibrium. Catch histories for some sectors were imputed from limited data. For the first time in a Bocaccio assessment, recreational catch was included as an input to the model. As in the previous work, this analysis indicates that Bocaccio exploitable stock biomass has declined significantly from the 1930s, with the steepest decline occurring from 1985 to 1995. The rate of decline slowed after 1995. While there is considerable uncertainty in estimating recent trends, there is no sign that the population has started to increase, and, more than likely, has continued to decline in the most recent decade. Based on the reference case results, the median estimate of stock size relative to its unfished stock size (B_{2012}/K) is 3.5%. The median estimate of current abundance relative to B_{msy} (biomass at maximum sustainable yield) is 7.0% with 90% confidence limits of 2.9% and 18.2% leaving little or no likelihood that the stock is currently above the lower Precautionary Approach reference point of $0.4*B_{msy}$, based on the reference case. Current harvests are approximately equal to the estimate of replacement yield. The impacts of alternative model assumptions from those used in the reference case were explored in 18 additional sensitivity runs but these results were similar to those of the reference case. Long term biomass projections were made for the reference case and a selection of the sensitivity runs over 5, 20, and 60 year scenarios under varying fixed harvest assumptions from 0-200 t/y. Results of the forecasts were presented relative to the DFO draft policy target reference points of $0.4*B_{msy}$ and $0.8*B_{msy}$. While the Bayesian approach used in this assessment provides a formal mechanism to include uncertainty in model output (including predictions), managers, and stakeholders are advised that not all sources of uncertainty have been addressed and that it is likely that the true uncertainty is even greater than that presented herein.

RÉSUMÉ

Le présent document contient une évaluation des stocks de bocaccio dans les eaux de la Colombie-Britannique, réalisée à partir de données de 2011. Les résultats du travail serviront aux gestionnaires et aux intervenants d'avis à court terme à propos de l'état actuel du stock et des effets probables des différents niveaux de prélèvement. Comme dans les travaux précédents, un modèle bayésien de production excédentaire a été employé. Il a été ajusté à partir d'un indice de biomasse dépendant de la pêche et de huit indices indépendants de l'indice de la biomasse, ainsi que des données de captures reconstituées de l'année 1935, à laquelle la population aurait été non exploitée et proche d'un équilibre naturel. Les données historiques sur les prises de certains secteurs ont été imputées à partir d'une information limitée. Pour la première fois dans une évaluation des stocks de bocaccio, la pêche récréative a été incluse dans les intrants du modèle. Comme le travail précédent, cette analyse montre que la biomasse du stock exploitable de bocaccio a considérablement diminué depuis les années 1930 et enregistré une chute brutale de 1985 à 1995. Le taux de déclin a ralenti après 1995. Malgré d'importantes incertitudes quant à l'estimation des tendances récentes, aucun signe n'indique un début de croissance de la population et cette dernière a, plus probablement, continué de décliner pendant la décennie passée. Compte tenu des résultats du scénario de référence, la médiane estimée de la taille du stock par rapport à la taille du stock non exploité (B_{2012}/K) est de 3,5 %. La médiane estimée de l'abondance relative actuelle par rapport à la B_{rms} (biomasse au rendement maximal soutenu) est de 7 %, avec des limites de confiance de 90 % à 2,9 % et 18,2 %, ce qui rend peu ou pas probable que le stock soit actuellement au-dessus du point de référence de l'approche de précaution, établi à $0,4*B_{rms}$ à partir du scénario de référence. Les captures actuelles sont à peu près égales au rendement de remplacement estimé. Les effets d'autres hypothèses du modèle que celles utilisées dans le scénario de référence ont été examinés dans 18 exécutions supplémentaires du modèle aux fins d'analyse de la sensibilité. Les résultats obtenus sont similaires à ceux du scénario de référence. Des projections à long terme de la biomasse ont été réalisées pour le scénario de référence et certaines des exécutions aux fins d'analyse de la sensibilité sur une période de 5, 20 et 60 ans, pour différentes hypothèses de prélèvement fixe variant de 0 à 200 tonnes/an. Les résultats des prévisions ont été présentés en comparaison des points de référence cibles de l'ébauche de politique du MPO, établis à $0,4*B_{rms}$ et $0,8*B_{rms}$. Bien que l'approche bayésienne utilisée dans le cadre de la présente évaluation offre un mécanisme formel pour inclure les incertitudes dans les résultats du modèle (y compris les prévisions), les gestionnaires et les intervenants doivent savoir que toutes les sources d'incertitude n'ont pas été étudiées et qu'il est probable que le degré d'incertitude réel soit plus important qu'il n'est indiqué dans le présent document.

1 INTRODUCTION

1.1 CONTEXT

In January, 2004, the Minister of the Environment received a document on Bocaccio (*Sebastes paucispinis*) from the Committee on the Status of Endangered Wildlife in Canada (COSEWIC). This document assessed the Bocaccio population in British Columbia (BC) waters as “threatened” (COSEWIC 2002). This prompted the Department of Fisheries and Oceans Canada (DFO) to conduct extended consultations with the government of BC, Aboriginal peoples, stakeholders, and the public on whether or not the Bocaccio population should be added to the List of Wildlife Species at Risk (Schedule 1) under the Species at Risk Act (SARA). Results of these consultations led the Governor in Council, through the Minister of the Environment, to refer the assessment back to COSEWIC in April 2006 for further information and consideration¹.

In December 2006, COSEWIC reaffirmed the original assessment without reassessing the species, citing an absence of new information that would lead to a change in the status of this species. In 2010, the Government of Canada, on the recommendation of the Minister of the Environment, acknowledged receipt of the COSEWIC Bocaccio assessment conducted under subsection 23(1) of SARA.

Following extensive review, the Governor in Council decided in 2011 not to add Bocaccio to the List of Wildlife Species at Risk. This decision was based on the recommendation of the Minister of the Environment and advice from the Minister of Fisheries and Oceans, taking into account the assessments provided by COSEWIC and those provided by DFO (DFO 2009a, Stanley et al. 2009a). It was determined that the costs of protection under SARA would likely outweigh the benefits to Canadians. However, the statement noted that protective measures would be taken under existing legislative tools such as the *Fisheries Act*, as well as non-legislative tools such as government programs and actions by non-governmental organizations.

Among the steps taken to provide protection to the Bocaccio population, an updated DFO assessment of Bocaccio was scheduled for 2011. In addition to updating the advice and enhancing the analysis, it would coincide with a COSEWIC re-assessment scheduled for 2011-2012.

1.2 OBJECTIVES

This assessment updates the previous Bocaccio assessment which used data current to 2007 (DFO 2009a, Stanley et al. 2009a). We refer readers to those documents for many of the background details on distribution, basic biology, data inputs, and modelling details. The basic objectives of this document are to:

- update the previous assessment with four more years of data (2008-2011);
- enhance the previous model;
- provide, with rationale, a Limit Reference Point, an Upper Stock Reference, guided by the DFO Sustainable Fisheries Framework (DFO 2009b);
- assess the status of the stock relative to the recommended reference points;
- predict the consequences of varying harvest levels on future population trends.

¹ Species at Risk Public Registry page for Bocaccio (including links to recommendations and decisions): http://www.sararegistry.gc.ca/species/speciesDetails_e.cfm?sid=740

The resulting information and advice may be used to assist development of groundfish management plans.

2 STOCK STRUCTURE

Consistent with previous documents, we treat Bocaccio as a single coastwide stock in BC waters. From the most recent US assessment of Bocaccio, Field 2011 notes that:

“from a population genetic perspective, all Bocaccio from British Columbia, Canada to Baja, Mexico, should probably be considered to be a single, panmictic unit”.

Buonaccorsi et al. (2012) note:

“We maintain that there is not enough evidence to reject the single homogeneous gene pool hypothesis for bocaccio rockfish... The lack of genetic divergence in this study does not support subdivision of the species’ range into separate management units, but does not necessarily refute division as moderate levels of population exchange may be sufficient to homogenize allele frequencies (e.g. Buonaccorsi et al. 2001), yet may not be sufficient for populations to quickly recolonize extirpated areas...”

...The findings of this study are concordant with an early allozyme study of bocaccio stock structure from southern to northern California which detected no significant allele frequency differences at two polymorphic loci (Wishard et al. 1980)...

We note that Bocaccio in the Puget Sound/Georgia Basin is considered by US sources as a "distinct population segment", based on life-history, environmental, ecological, and genetic information (Drake et al. 2010). The genetics conclusion is based on observations on “sister” species (Alexandra Valentin, COSEWIC, personal communication, 2012).

Field et al. (2009) note in the 2009 US assessment that:

“The National Marine Fisheries Service (NMFS) recently issued a proposed rule (and request for comment) to list the population of Bocaccio in the Georgia Basin (Puget Sound, Washington and the Strait of Georgia in BC as endangered (at high risk of extinction) under the Endangered Species Act (ESA). This proposed rule came about as a result of a petition to enlist this and several other population units of rockfish in this region (the other four species were canary, yelloweye, greenstriped and redstripe rockfish). Of these five only bocaccio is proposed to be listed as endangered...”

The proposed rule is based on the evaluation of abundance trends, spatial structure of the populations, and the suite of somewhat unique threats in this ecosystem. Among the factors related directly to bocaccio are the rapid decline and current total absence of bocaccio in recreational rockfish catches within the Georgia Basin (consistent with a substantial overall decline in the catch rates of all rockfish, but of a greater magnitude), the highly variable nature of bocaccio recruitment, and the observation that historical length composition data were indicative of multiple strong cohorts (interpreted as evidence that fish present in the ecosystem were unlikely to be infrequent strays from the coastal population).”

Our interpretation of the COSEWIC guidelines for recognizing Designatable Units (Guidelines for Recognizing Designatable Units: http://www.cosewic.gc.ca/eng/sct2/sct2_5_e.cfm) is that the US work provides only a partial justification for assuming a separate Puget Sound/Georgia Basin population. Certainly no genetics research work has been conducted on this issue, nor are there any biological data for the BC portion of this area for comparing life history parameters.

Presumably, the virtual absence of catch and sample data as well as the lack of relative abundance indices for the BC portion of the Puget Sound/Georgia Basin would render such an assessment “data limited”. However, given the possibility that there is a separate population and the sparseness of the data from this area, we exclude these data from the analysis as in the previous assessment. No separate assessment has been conducted on the BC portion of the Puget Sound/Georgia Basin Bocaccio population.

3 DISTRIBUTION

The distribution of Bocaccio catch observations in BC waters is provided in Figure 1. This figure shows the location of capture for all observations of Bocaccio in the most recent 10 years (2002-2011). We did not attempt to examine whether distribution has changed over time. The commercial data are not comparable over time and the time series of the various surveys are relatively short (see below).




Figure 1. Area of occupancy for Bocaccio based on survey and commercial observations from 2002-2011 (commercial trawl 2002-2011; commercial HL 2006-2011). Figure indicates all 2 km x 2 km cells with at least one record of Bocaccio capture.

4 LIFE HISTORY PARAMETERS

4.1 SAMPLE SOURCES

The estimates of size-at-age and maturity-at-age were derived from 1,212 aged specimens collected between 2001-2010; these included 940 used in the previous assessment (DFO Groundfish GFBio database). These samples came from both research survey and commercial fishery catches. The commercial samples were obtained from at-sea observer and port samples of both midwater and bottom trawl catches. As noted in Stanley et al. 2009a, we concluded that there were too few data to explore the influence of catch source, gear, location, depth, and season on the estimates of size-at-age, maturity-at-age, or length/weight. An updated summary of the age composition is provided in Figure 2.

There are more males than females in the age-length samples for all ages, but it is not known whether this reflects higher natural mortality, higher fishing mortality, or a combination of both. It may also represent different selectivities between sexes in these samples which were collected from surveys and commercial fishing since 2001.




Figure 2. Bubble plot of proportions-at-age for female and male Bocaccio for 2001-2010, all sources combined. Area of the bubble is proportional to proportion-at-age within each sex. Youngest age showing in 2009 and 2010 corresponds to 3-year-olds. No age data are available from prior to 2001.

4.2 GROWTH, AGE-AT-MATURITY, AND NATURAL MORTALITY

Growth was estimated using same methods as before (Stanley et al. 2009a), using the updated data. A von Bertalanffy growth model was fitted to the length-age observations (Figure 3 and Table 1). The estimates of k and L_{inf} are precisely determined owing to the fairly large sample sizes. The estimate of t_0 is poorly determined because there was only one observation below age seven for males and three observations below age six for females.

The estimates of maturity-at-age for female Bocaccio were updated with a subset of the additional data ($n=321$)² (Figure 4). However, we used a different method in which proportion mature at age was estimated by applying a cumulative, renormalized lognormal density function. The age at which the maturity-at-age function intersects with zero mature was set at the oldest immature age at which zero animals were found to be mature. The number found to be mature in each set of samples at each age was modeled to be a binomial random variable with the probability predicted by the renormalized cumulative lognormal density function. The parameters estimated included the median value and standard deviation in the natural logarithm of age for the lognormal density function (see King et al. 2012 for details on the methodology). The median estimate of age at 50% maturity for females was estimated to be 7.1 years as opposed to the 8.5 years estimated previously (Stanley et. al. 2009a). The advantage of applying the cumulative renormalized lognormal function over the logistic function is that it provides a much better fit to the maturity data for Bocaccio. The best fitting logistic function (not shown) markedly under-predicted the observed positive fraction of mature values for the youngest ages.

Table 1. Updated growth parameters for Bocaccio.

| Sex | Parameter | 2012 | | 2009 |
|---------|-------------------|----------|---------|----------|
| | | Mean | SD | Mean |
| Females | L_{inf} (cm) | 79.520 | 0.630 | 78.32 |
| | k (yr^{-1}) | 0.162 | 0.002 | 0.163 |
| | t_0 (yr) | -0.510 | 0.380 | -1.20 |
| Males | L_{inf} (cm) | 69.180 | 0.150 | 69.98 |
| | k (yr^{-1}) | 0.177 | 0.001 | 0.108 |
| | t_0 (yr) | -1.970 | 0.400 | -8.46 |
| Females | a | 8.57E-09 | 1.0E-09 | 3.58E-05 |
| | b | 3.10 | 0.028 | 2.754 |

² As in previous work, we used observations on female maturity from months of February-July owing to difficulty in field-staging of rockfish maturity (see Stanley and Kronlund, 2004 for an explanation of the methodology).





Figure 3. Observed and estimated length-at-age for a. female and b. male Bocaccio.




Figure 4. Observed and estimated proportion mature-at-age for female Bocaccio. The median refers to the age at which 50% of females are predicted to be mature.

A graphical comparison of the proportion mature-at-age with the age frequency samples from the commercial fishery implies that recruitment to the fishery and the maturity ogives are similar, which may indicate that there is limited exploitation on juvenile females (Figure 5).




Figure 5. Female Bocaccio proportion mature-at-age (line) compared with histogram of age frequency of females, for all samples combined.

As in Stanley et al. 2009a, we treated instantaneous rate of natural mortality for females, M , as a lognormal random variable with a median of 0.075 yr^{-1} and standard deviation in log space equal to 0.25. Generation time is treated as approximately 20 years.

5 CURRENT MANAGEMENT AND MONITORING OF BOCACCIO CATCHES

Bocaccio are captured incidentally in all commercial groundfish trawl and most hook-and-line fisheries, as well as the salmon troll and recreational. There is currently no directed fishing for Bocaccio in any fishery. Some targeting took place during earlier decades of the groundfish trawl fishery.

There are currently no regulations that specifically limit trawl catches of Bocaccio although landings of non-quota rockfish in the trawl fleet (including Bocaccio), are limited to a maximum of 15,000 lbs per trip, all non-quota species combined³. However, to address the concerns for Bocaccio, a voluntary program for the trawl fleet was developed and implemented in 2004 in which groundfish trawl vessels relinquished all landed Bocaccio catches and directed the proceeds for research and management purposes. These actions resulted in an approximate halving of Bocaccio trawl landings after 2004 relative to level of landings prior to that year (Figure 6, Appendix Table 1 and Appendix Table 2). This voluntary relinquishment program remains in place at this time.

In the commercial groundfish hook-and-line and trap fisheries (HL fisheries), Bocaccio is managed as part of an aggregate of "other" rockfish which is applied to non-quota rockfish. For Inside ZN rockfish fishing (i.e., inside waters of Vancouver Island), the combined catch in a trip of "other" rockfish must be less than or equal to the combined catch of Yelloweye, Quillback, Copper, China, and Tiger Rockfish. For HL fishing in outside waters, there is a trip limit of 5,000 pounds for non-quota rockfish combined. Recreational catches are constrained by "rockfish" daily bag limits of 0 to 5, depending on the area.

³ See the 2011-2013 Integrated Fishery Management Plan for Groundfish at http://www.pac.dfo-mpo.gc.ca/fm-gp/mplans/ground-fond_2012-13.pdf

Trawl catches (not including those from inside waters) have been monitored since 1991 with independent 100% dockside coverage of landings and, since 1996, with 100% observer coverage of at sea catches. Total catches from this fishery are considered accurate since 1996.

Catches in the groundfish HL fishery have been monitored with 100% independent dockside monitoring for all sub-sectors since 1996 (ZN rockfish in 1995) and with a 100% electronic monitoring of catches at sea since 2006. Catches for this fishery are considered accurate since 2006 (see Stanley et al. 2009b).

Rockfish catches in the recreational fishery, depending on the area, are monitored primarily through a combination of creel surveys, aerial flights to estimate effort, and harvester logs. The program is primarily designed to estimate the catches of salmonids, but there has been a concerted effort to improve the monitoring of groundfish species in recent years.

6 COMMERCIAL CATCH DATA

This assessment uses catch data from the same fisheries as were modelled in the previous assessment, with four additional years of observations (2007/2008-2010/2011). This assessment also models the recreational catches for the first time in a Bocaccio assessment (Table 2).

Table 2. The seven fisheries modelled in this stock assessment, showing the years these fisheries were included. Catches from four of these fisheries were assumed known without error ("Fixed"). Catches from the remaining fisheries were estimated as described in Stanley et al. 2009a and below.

| Gear | Sector | Years | Fixed or Estimated |
|-------------|---------------------|--------------|---------------------------|
| Trawl | US domestic | 1935-1980 | Fixed |
| Trawl | CDN domestic | 1950-2011 | Fixed |
| Trawl | Soviet and Japanese | 1965-1977 | Fixed |
| HL | CDN Rockfish ZN | 1940-2011 | Fixed |
| HL | CDN and US Halibut | 1935-2011 | Estimated |
| Troll | CDN Salmon troll | 1935-2011 | Estimated |
| Handline | Recreational | 1935-2011 | Estimated |

In this document, "catch" refers to total removals by fishing gear, summing both the retained (landed) catch and the discarded catch. We assume that all Bocaccio die after capture, so this sum is equivalent to total fishery-generated mortality. Bocaccio have been predominantly a non-directed (or bycatch) species in all BC fisheries.

6.1 TRAWL CATCH

Catch for all trawl fisheries were input as fixed values, known without error (Figure 6, (Appendix Table 1 and Appendix Table 2). Details on how historical catches were reconstructed are provided in Stanley et al 2009a. Data were updated for 2008-2011. Catch estimates for 2012 were assumed to be equal to 2011.




Figure 6. Domestic (U.S. and Can.) and foreign (Soviet, Japanese, and Polish) trawl landings (1930 to 2011).

6.2 HOOK-AND-LINE CATCH

As in the previous assessment, we divided the HL and trap fisheries into three sectors: rockfish ZN (set-line, and handline and lingcod troll), halibut (set-line), and salmon troll. Catches for the rockfish ZN fishery, which primarily targets rockfish and lingcod, were taken directly from DFO catch databases as fixed values, known without error (Figure 6, Appendix Table 1 and Appendix Table 2).

Catches of Bocaccio in the halibut, salmon troll, and recreational fisheries were estimated with the same methodology described in the previous assessment for halibut and salmon troll (Appendix Table 1, Appendix Table 2, and Appendix Table 3). The time series of halibut catch is estimated as a function of fishing effort in outside waters. This effort series was updated for this assessment with an additional four years, ending in 2011 (Figure 7).

The model predicted annual catches in the halibut fishery from the observed halibut effort ($E_{f,y}$) (see Eq. B1 in Stanley et al. 2009a, Appendix B). The catchability coefficient, k_f , was estimated from the observed halibut catches during the years 2006-2011, with a non-informative prior for k_f (Stanley et al. 2009a) and with updated catch records ending in 2011. The imputed values of catch for these fisheries are provided below in the section on model results (Section 11.4 and Figure 21). The salmon troll effort series was extended up to 2010; 2011 data were not available at the time of report preparation. Effort values for 2011 and 2012 were assumed to be equal to 2010.




Figure 7. Time series of annual effort used for imputation of total catch in the salmon troll, halibut, and recreational fisheries.

6.3 RECREATIONAL CATCH

Earlier Bocaccio assessments treated recreational catches as negligible relative to other fisheries and were not included in the analyses. However, the relative importance of this fishery has increased, given the reduced catches in the trawl fishery and the low estimates of current biomass. This means that the impact of recreational exploitation rates needs also to be considered.

Similar arguments could be made for the inclusion of exploitation of catches of Bocaccio by other fisheries (e.g., salmon seine and gillnet, prawn trap and First Nations); however, this was not attempted for this assessment. The exploitation by each of these fisheries is assumed to be lower than that for the recreational fishery, although considered collectively, there may be having an impact. However, the lack of available data precludes the sensible inclusion of these fisheries.

Consistent with the rest of the assessment, we did not include recreational catch from the inside waters of Vancouver Island (Strait of Georgia, Juan de Fuca Strait, and Queen Charlotte Strait), although catches of Bocaccio have been reported from Area 12, Queen Charlotte Strait (Table 3). The appearance of Bocaccio in 2008 in Table 3 reflects ongoing effort to improve monitoring of non-salmonids in this fishery.

Table 4 summarizes available recreational catch estimates of Bocaccio from outside waters of the BC. South Coast data exclude catches and effort from waters inside of Vancouver Island (i.e., Area 20-Juan de Fuca Strait, the Strait of Georgia, and Areas 11 and 12-Queen Charlotte Strait). Note that although effort units are collected as angler days in the Central Coast data, they were treated as “boat days” in the discussion below. North coast estimates were only available for “Rockfish”, so we converted these to pieces of Bocaccio by using the observed yearly bycatch ratio (Bocaccio/Rockfish) for the South and Central Coasts combined.

Table 3. Estimates of Bocaccio catch from Minor Area 12 (Inside waters of Vancouver Island – north) (DFO, unpublished data).

| Year | Minor Area 12 | | |
|------|----------------------|----------------------|------------|
| | Rockfish (pieces) | Bocaccio (pieces) | Boat Trips |
| 2000 | 11136 | | 18083 |
| 2001 | 7670 | | 10825 |
| 2002 | 1415 | | 5016 |
| 2003 | 4918 | | 14059 |
| 2004 | 5924 | | 16367 |
| 2005 | 4762 | | 18682 |
| 2006 | 9370 | | 15901 |
| 2007 | 5407 | | 16737 |
| 2008 | 7028 | 127 | 12914 |
| 2009 | 9827 | 107 | 15080 |
| 2010 | 9335 | 41 | 14981 |
| 2011 | 8298 | 11 | 15623 |

Assuming that Bocaccio catches were underestimated for the north coast in 2007 (3 pieces), we used the 2008-2010 data for all areas. This indicated a mean annual reported catch of 250 pieces coastwide. This equates to about 1 t using the mean weight of 4.3 kg observed in commercial fisheries. Assuming a recent coastwide fishing effort of about 100,000 days (including an extra 14,000 days for the North Coast), this catch rate would correspond to about 0.0025 Bocaccio/day coastwide. This implies a Bocaccio/Rockfish catch ratio of 0.66%, which is consistent with the ratios shown in Table 4.

We used simple methods to combine recreational data sources by area; however, the intent was only to develop a starting minimum assumption of coastwide catch. The resulting value of 1 t per year was so low relative to other fisheries we suggest no additional work is warranted.

Table 4. Estimates of recreational catch of Bocaccio 2000-2010 (unpublished DFO data). Note that “total boat days” does not include a north coast estimate. Values in italics were estimated from monitoring data.

| Year | Total | | | South Coast (excl. SoG) | | | Central coast and Area 11 | | | North coast | | |
|------|--------------------|------------------|-----------|-------------------------|------------------|-----------|---------------------------|------------------|----------------|--------------------|--|----------------------------|
| | Rockf. (pieces) | Boc. (pieces) | Boat days | Rockf. (pieces) | Boc. (pieces) | Boat days | Rockf. (pieces) | Boc. (pieces) | Angler Days | Rockf. (pieces) | Catch Ratio from SC, Area 11 and CC combined | Boc. (pieces) (est.) |
| 2000 | 34093 | 0 | 55788 | 14100 | | 55788 | 0 | 0 | 0 | 19993 | | |
| 2001 | 39481 | 0 | 60811 | 21048 | | 60811 | 0 | 0 | 0 | 18433 | | |
| 2002 | 50466 | 0 | 102511 | 30437 | | 75262 | 4603 | 0 | 27249 | 15426 | | |
| 2003 | 33386 | 2 | 113384 | 21965 | | 84124 | 4234 | 2 | 29260 | 7187 | | |
| 2004 | 42516 | 4 | 115518 | 29620 | | 81825 | 4952 | 4 | 33693 | 7944 | | |
| 2005 | 55601 | 0 | 112316 | 43157 | | 79202 | 4713 | 0 | 33114 | 7731 | | |
| 2006 | 52129 | 14 | 114891 | 37293 | | 79401 | 7556 | 14 | 35490 | 7280 | | |
| 2007 | 44503 | 87 | 99082 | 32460 | 78 | 64632 | 10422 | 9 | 34447 | 1622 | <i>0.0020</i> | <i>3</i> |
| 2008 | 69653 | 279 | 95184 | 39829 | 243 | 66908 | 8798 | 36 | 28155 | 21026 | <i>0.0057</i> | <i>120</i> |
| 2009 | 62127 | 261 | 93027 | 31452 | 220 | 71768 | 7238 | 41 | 21101 | 23437 | <i>0.0067</i> | <i>158</i> |
| 2010 | 65379 | 212 | 86580 | 30035 | 117 | 66208 | 7076 | 95 | 20210 | 28268 | <i>0.0057</i> | <i>162</i> |

For comparison, the commercial HL (non-trap and non-dogfish) fisheries catch about 0.31 Bocaccio/day (2006-2011), or 100 times greater and the catch ratio (Bocaccio/Rockfish) is about 1% (50% higher).

While it is obvious that monitoring has improved in recent years, we suggest the true recreational catch is likely underestimated. This can be inferred from the large number of unidentified “Rockfish”. Therefore, we suggest that 1 t/y is suitable as a minimum current (2010) coastwide estimate of the recreational catch of Bocaccio. There is no analytical basis for deriving a maximum estimate but we proposed a theoretical upper limit of about 10 t/y (about 2,300 pieces/y). This translates to a catch rate of 0.023 Bocaccio/day, or 1 fish for every 43 boat days of sport fishing (100,000 days/ (10 t/4.3 kg)).

The historical recreational catch of Bocaccio in outside waters was estimated using the same methodology as in the halibut and salmon troll fisheries. We used the outside waters recreational fishing effort series, compiled for the 2011 Quillback Rockfish assessment (Yamanaka et al. 2012) (Figure 7 and Appendix Table 3). Data for 2011 were not available at the time of report preparation; consequently, recreational effort for 2011 and 2012 was assumed to be the same as that in 2010.

As indicated above, a single approximation of the bycatch of Bocaccio in recreational fisheries was formulated for the year 2010. This was originally specified as a triangular distribution with a minimum of 1 ton, a peak at 5 tons and a maximum at 10 tons (Figure 8). This prior was treated as a pseudo data point just as the one for the average catch in outside waters salmon troll fisheries had been treated as a pseudo data point. To allow for improved numerical performance in the Bayesian model, the triangular distribution was replaced by a truncated normal density function for the estimated catch in 2010, with the minimum value set to 1.0 ton. The closest fit between the normal and initial triangular distribution was obtained by setting the mean of the normal density function (without truncation) to 5.2 tons and CV to 0.4 (Figure 8). The probability density of the data point was computed using the model predicted value for the recreational catch in year 2010 as follows:

$$C_{r,2010} \sim Normal\left(\hat{C}_{r,2010}, \left(CV \times \hat{C}_{r,2010}\right)^2\right)$$

where $C_{r,2010}$ is the data point for the recreational catch in 2010 and $\hat{C}_{r,2010}$ is the model-predicted recreational catch in 2010. This is predicted in the same way that the halibut and troll catch is predicted (i.e., by assuming that the fishing mortality rate from recreational fishing is directly proportional to the annual recreational fishing effort):

$$\hat{C}_{r,2010} = B_{2010} \left(1 - \exp\left(-k_r E_{r,2010}\right)\right)$$

where B_{2010} is the model predicted biomass in 2010, k_r is the recreational catchability coefficient, and $E_{r,2010}$ is the recreational fishing effort in 2010. The estimated recreational catch by year is provided below in model output (Section 11.4 and Figure 21, Appendix Table 1, Appendix Table 2 and Appendix Table 3).

Note that the above process, which bases catchability on 2010 results, leads to estimates of higher catches going backwards over time, reaching a peak median estimate of 33 t in 1984 then slowly declining to near 0 by 1945. This trend simply reflects the effect of greater Bocaccio abundance in combination with the trend in recreational effort. These estimates are highly uncertain (Figure 21), dependent on the assumed prior based on the 2010 estimated catch (Figure 8) and the additional assumption that catchability has been constant in time (e.g., constant fishing behaviour and gear). While it is extremely unlikely that catchability has been constant during the past 60 years for these effort-driven fisheries (recreational, salmon troll, and

halibut fisheries), we lack any basis for estimating changes in catchability for these fisheries and feel that this methodology is a reasonable means for accounting for the impact of these fisheries.

In the case of recreational fishery, annual current catches which lie between 1 and 10 t seem to be a reasonable range; however, there is little basis to defend the modal choice of 5 t, other than being the approximate midpoint. The true mean could be as low as 2-3 t, which would reduce the historical series by half. However, the mean could be closer to 10 t, leading to the opposite effect.




Figure 8. Plot of the initial prior distribution for recreational catch of Bocaccio in BC in 2010 from a qualitative interpretation based on recreational catch data and interviews. A normal approximation of the prior is also shown. This latter distribution was used to compute the probability of this prior approximation of recreational catch given model predicted catches of Bocaccio in 2010.

7 ABUNDANCE TRENDS - COMMERCIAL CPUE

We included the same commercial bottom trawl CPUE index for 1996/1997 to 2003/2004 used previously (Stanley et al. 2009a) as an index of abundance in the assessment model. This index was based on commercial catch and effort data collected from bottom trawl fishing by independent observers over the period 1996–2004. As explained in Stanley et al. (2009a), we did not use catch and effort data prior to 1996 because these data are neither trustworthy nor were they collected and archived in a comparable fashion.

We only used data through to 31 March 2004, which was the end of the “fishing year” at that time. After this date, in response to concerns expressed about the status of Bocaccio, as noted earlier, most participants in the trawl fishery voluntarily agreed to relinquish⁴ the value of all Bocaccio landings. This initiative not only removed the incentive to target Bocaccio, but also encouraged harvesters to avoid Bocaccio. Trawl catches in this sector declined from around 200-250 t annually to nearly 100 t per year by the 2006/2007 fishing year. Consequently, we believe that Bocaccio catch rates after the 2003/2004 fishing year are not comparable with the earlier period and ended the series in 2003/2004. The standardized and nominal trends

⁴ The individual harvester did not receive any payment for Bocaccio landings. Revenue was “relinquished” to the Canadian Groundfish Research and Conservation Society.

indicate there was little change in CPUE from 1996/1997 to 2003/2004 (Figure 9 and Appendix Table 4).




Figure 9. Comparison of the lognormal and binomial standardized commercial trawl CPUE indices for Bocaccio. The error bars show \pm 95% confidence bounds.

Catch data were standardized using Generalized Linear modelling (GLM) methods (Stanley et al. 2009a). The nominal and standardized indices, as well as other treatments of the data, provided similar trends over the selected time period, probably indicating that fishing practices changed little over this period. This stability across treatments does not validate the series as an index of abundance, but indicates that alternative methods are unlikely to provide a significantly different signal.

Concerns over comparability in fishing behaviour have led us to exclude the use of commercial catch rates from the groundfish HL fisheries. Improved monitoring, starting in 2006, has provided accurate estimates of catch (landings and discards) of Bocaccio from that year forward. However, we think it is likely that fishing behaviour has changed and continues to change as the fleet adapts to Individual Vessel Quotas (introduced in 2006), thus rendering it unlikely catch rates in this fishery will provide usable indices of relative abundance for Bocaccio.

8 ABUNDANCE TRENDS - SURVEYS

We used the results from eight surveys in this stock assessment (Figure 10 and Figure 11, Table 5 - Table 7). Results for the two shrimp surveys, and the four Groundfish synoptic surveys were updated with 2008-2011 results using the methodology described in Stanley et al. (2009a) (Figure 12 and Figure 13, Appendix Table 5 - Appendix Table 10). We again used biomass indices from the US NMFS Triennial survey which ended surveying in BC waters in 2001 (Figure 12 and Appendix Table 11).

We included results from the West Coast Haida Gwaii Groundfish Synoptic Survey (WCHG) for the first time (Figure 13 and Appendix Table 7). This survey was excluded previously because there were only two data points. Abundance indices for this survey were derived in the same manner as followed for the other synoptic trawl surveys (see Stanley et al. 2009a).

We also included the International Pacific Halibut Commission (IPHC) longline survey results for the first time (Figure 12 and Appendix Table 12). The method of calculating the index for this longline surveys is provided in Section 21, Appendix B. The IPHC survey is a fixed station longline survey of approximately 170 stations. Although initiated before 2003, we have only used results from 2003-2011 because this was the first year that groundfish catch (in pieces)




Figure 10. Locations of Shrimp trawl, US NMFS Triennial bottom trawl, and IPHC HL longline surveys.




Figure 11. Locations of Groundfish Synoptic bottom trawl surveys.




Figure 12. Plots of biomass estimates for Bocaccio from the a. West Coast Vancouver Island shrimp trawl, b. Queen Charlotte Sound shrimp trawl, c. U.S. Triennial survey, and d. IPHC longline surveys. Bias corrected 95% confidence intervals from 1,000 bootstrap replicates are plotted.




Figure 13. Plots of biomass estimates for Bocaccio from the a. West Coast Haida Gwaii; b. Hecate Strait, c. Queen Charlotte Sound and, d. West Coast Vancouver Island Groundfish Synoptic trawl surveys for 2003 to 2011. Bias corrected 95% confidence intervals from 1,000 bootstrap replicates are plotted.

9 TRAWL SURVEY CATCHABILITY

As in the 2009 assessment, an informative prior for trawl survey catchability, q , was applied to improve the precision in estimates about stock biomass (Stanley et al. 2009a, McAllister et al. 2010). In this assessment, one additional trawl survey index of abundance for from the WCHG survey was included so the informative prior for q was extended to include this survey (Table 8) (see Appendix Figure 1 and Appendix Table 13 for the confidence limits and marginal density functions for q -gross).

Results from a rockfish gillnetting experiment (Matthews et al. 1989) were also compiled to provide an estimate of the ratio of the density of Bocaccio in untrawlable and trawlable areas. The value for the ratio of the mean catch rate means across 21 sets in untrawlable and 20 sets in trawlable areas was 1.44. The lognormal standard deviation of the ratio was 0.28. A lognormal density function was used to compute the probability of the experimental observation given the q model predictions of the ratio (McAllister et al. 2010). The updated prior for the ratio of catch rate between untrawlable and trawlable areas caused the prior median values for q to increase slightly in value and for the joint prior distribution for q to become slightly more precise than that used in the 2009 assessment. The higher q is associated with an overall downward scaling of biomass.

The joint prior for survey catchability was approximated by a seven dimensional lognormal density function (one dimension for each of the seven trawl survey datasets) incorporating the median and covariance of $\log q$ which were used as the q prior distribution in the assessment model.

Table 8. Posterior means and CVs for q -gross ($qgfin$) for the current and 2009 assessments

| Survey | 2012 | | 2009 | |
|------------------------------|--------|------|--------|------|
| | Mean | CV | Mean | CV |
| #1 - WCVI groundfish | 0.1110 | 0.69 | 0.0703 | 0.77 |
| #2 - QCSd-groundfish | 0.0720 | 0.72 | 0.0459 | 0.80 |
| #3 - HS - groundfish | 0.0105 | 0.76 | 0.0067 | 0.83 |
| #4 - WCHG - groundfish | 0.0034 | 0.71 | 0.0021 | 0.79 |
| #5 - WCVI Shrimp | 0.0048 | 1.40 | 0.0030 | 1.46 |
| #6 - QCSd Shrimp | 0.0005 | 2.64 | 0.0004 | 2.74 |
| #7 - US Triennial groundfish | 0.0605 | 1.56 | 0.0474 | 1.73 |

10 BAYESIAN SURPLUS PRODUCTION MODEL

This assessment used the non-equilibrium, age-aggregated Bayesian surplus production (BSP) model described in Stanley et al. (2009a). It is a state-space version which incorporates stochastic process error in the fish stock dynamics (Meyer and Millar 1999) and thereby permits a more thorough accounting of uncertainty in estimates of stock biomass, stock projections, and deviations as compared to a deterministic surplus production model. A Bayesian statistical approach was adopted to fit the model to data, allowing for the use of informed priors which incorporated information and expert judgements. The BSP model was fitted to eight sets of survey abundance indices and the one commercial CPUE series to reconstruct historical trends in abundance of Bocaccio. The fitted model was then used to evaluate the future trends in abundance based on alternative total allowable catch (TAC) policies. TAC refers to total combined catch from all modelled fisheries, including recreational catch for the first time.

We use a version of the Schaefer surplus production function (Hilborn and Walters 1992) that applies continuous fishing mortality rate equations (Prager 1994, and see Stanley et al. 2009a):

$$\text{Eq. 1} \quad B_t = B_{t-1} + B_{t-1}r\left(1 - \frac{B_{t-1}}{K}\right) - F_{t-1}B_{t-1}$$

where B_t is stock biomass in year t , r is the maximum intrinsic rate of increase, K (or B_0), is the average unfished stock size or carrying capacity, and F_t is the instantaneous fishing mortality rate in year t . The estimation performance of a Bayesian version of this model was evaluated and found to perform acceptably under a range of conditions using simulation testing of the state space version of this model (unpublished work by first and second author). Earlier simulation testing of a deterministic version also produced acceptable performance (McAllister and Kirkwood 1998). This testing included misspecification of the priors, as long as the priors for key parameters (e.g., r and constants of proportionality for stock trend indices, q) were not overly precise or strongly biased (McAllister and Kirkwood 1998). The version used in this assessment provides more accurate representations of fish stock dynamics than a deterministic version or discrete harvest rate version, especially when fishing mortality occurs throughout the year and when exploitation rates are high. It is slightly more cumbersome because the annual fishing mortality rate (F_t) must be solved numerically rather than analytically as in the discrete version (see McAllister and Babcock 2002 and McAllister et al. 1999; 2001a for additional details on the model).

We applied a state-space version of the BSP that incorporates lognormal deviates from total annual biomass predictions:

$$\text{Eq. 2} \quad B_t = \left(B_{t-1} + B_{t-1}r\left(1 - \frac{B_{t-1}}{K}\right) - F_{t-1}B_{t-1} \right) \exp\left(\varepsilon_t - \frac{\sigma_p^2}{2} \right)$$

where the prior probability distribution for the process error term is given by

$\varepsilon_t \sim \text{Normal}(0, \sigma_p^2)$. Values for ε_t from 1935 to 2011 were treated as estimated parameters and σ_p was set at 0.1. In the 2009 assessment, sensitivity tests showed that stock status and projection results were insensitive to two alternative settings of 0.05 and 0.15 for σ_p .

Consequently, these sensitivity runs were not repeated in this assessment. No attempt was made to estimate the process error variance or the observation error variance, owing to the paucity of time series data that could inform estimates of variance in ε_t and the low precision in most of the indices.

The reference case prior distributions for K , r , the ratio of stock size in 1935 to K (B_{1935}/K), and the constants of proportionality (q) for the stock trend indices are provided in (Table 9) (see Appendix G of Stanley et al. 2009a for the methodology used to develop these priors). As was done in 2009, the prior for the maximum intrinsic rate of increase, r , was developed using a demographic approach (McAllister *et al.* 2001b). This approach was based on available life history data on growth, the natural mortality rate (M), maturity-at-age, and the Ricker stock-recruit steepness parameter, developed from a hierarchical meta-analysis of rockfish stock-recruit data (Forrest et al. 2010). The posterior predictive distribution for the Ricker steepness from Forrest et al. (2010) was approximated using a transformed beta density function with minimum of 0.2, mean of 0.93, and standard deviation of 0.42.

The method used to develop the prior for r in the 2009 assessment only accounted for uncertainty in M and steepness. A similar methodology was applied in this assessment, except that the CV of the M -prior was increased from 0.20 to 0.25 yr^{-1} . In addition, the methodology was expanded to include empirical uncertainty in the parameter estimates for growth, the length-weight relationship, the proportion mature at age, and the Ricker steepness parameter (from Forrest et al. 2010 and see Yamanaka et al. 2012). A Ricker stock-recruit function was adopted in preference to the Beverton-Holt stock formulation because there has been a report of cannibalism in Bocaccio (Love et al. 2002). The prior for r was then developed from a simulation model which included these life history parameters, represented as priors by their posterior mean and covariance matrix (see Eq. 26 to Eq. 32 in Appendix G, Stanley et al. 2009a). The mean and standard deviation (SD) for the r -prior used in this assessment were 0.1067 and 0.039 (Figure 14), which are similar to the mean of 0.117 and SD of 0.035 used in the 2009 assessment. The prior distribution for r is approximated in the model by using this mean and SD to describe a normal distribution. Prior distributions for r , representing higher and lower levels of productivity, were developed in the same manner for use in model sensitivity runs (Figure 14).

Table 9. Prior pdfs of parameters K , q for the commercial CPUE data, P_0 , and r .

| Parameter | Prior Density function | Comments |
|--|-----------------------------------|--|
| K | Uniform (500, 200,000) | Units in tons |
| q for commercial cpue and the IPHC index | Proportional to $1/q$ | This prior is non-informative with respect to K and stock biomass (See Stanley et al. 2009a: Table 4, Appendix F11 and F13 for key details on the informative prior for the survey q_s). |
| P_0 | Lognormal($\ln(0.9)$, 0.2^2) | This indicates that the stock was near to carrying capacity in 1935. |
| r | Normal(0.1067 , 0.0391^2) | The relatively low prior mean comes largely from the late median age at maturity of 7 years. It also comes from the relatively low estimates of recruits per ton of spawner biomass at the origin of the stock-recruit function which in turn derives partly from the low prior mean for steepness obtained from the meta-analysis of rockfish stock recruit data (Forrest et al. 2010). |




Figure 14. Fitted normal prior density function of the maximum intrinsic rate of increase, r , for BC Bocaccio. The square dots show the frequency distribution of values simulated from the stochastic demographic model for r . The 2009 reference case and 2012 reference or base case, and low and high prior r cases, are shown.

The Schaefer surplus production model assumes that B_{msy}/K (or B_{msy}/B_0) occurs at 50% of K . This is a property of the model parameterisation and does not reflect the productivity or other biological characteristics of Bocaccio. However, we have chosen this parameterisation for our reference case because we believe that this ratio is a credible representation of Bocaccio for a number of reasons. One reason is that a recent hierarchical meta-analysis of stock-recruit data for rockfish (Forrest et al. 2010) indicated that the credible range for the median B_{msy}/K by

species spanned 0.15 to 0.5 for the Beverton-Holt stock-recruit function and 0.35 to 0.5 when the Ricker stock-recruit function was fitted to the same data (Figure 15). Furthermore, when the steepness parameter for either stock-recruit function approaches lower values, which tend to be more consistent with the low value for r estimated for Bocaccio, the estimated value for B_{msy}/B_0 tends to approach 0.5 (Figure 15) (Forrest et al. 2010).

Nevertheless, recognising that the choice of $B_{msy}/K=50\%$ was arbitrary; we investigated three alternative forms of the surplus production function as sensitivity tests, wherein we fixed the B_{msy}/K ratio at 0.3, 0.4, and 0.6. We used a variant of the Fletcher generalized surplus production function (Quinn and Deriso 1999), which allowed the value of B_{msy}/K to take on any value between 0 and 1. We use this form because the classical forms of the Pella-Tomlinson and Fletcher generalized surplus production functions have the property where r and the value for B_{msy}/K are negatively correlated, with r becoming infinity when B_{msy}/K decreases below the value of $1/e$ (~ 0.37) (Quinn and Deriso 1999). The variant employed in these sensitivities uses a parabolic Schaefer production form for the portion of the production function below B_{msy}/K , thus allowing the Schaefer production function to be continuous with the Fletcher form at MSY (McAllister et al. 1999). This also permits the prior for r to be incorporated directly into the generalized model, which is not permitted in the classic generalized form. The corresponding Fletcher functions (dotted curves in Figure 16) are shown for B_{msy}/K implementations of 0.3 and 0.4. Note that the Schaefer model is a special case of the Fletcher model, when B_{msy}/K is fixed at 0.5.





Figure 15. Estimated values for B_{msy}/B_0 under the Ricker and Beverton-Holt stock-recruit models as a function of the steep parameter in the a) Ricker and b) Beverton-Holt stock-recruit functions (Forrest et al. 2010).




Figure 16. A plot of the reference case Schaefer and three alternative production functions applied in evaluations of the sensitivity of results to different model settings. All plotted production functions are referenced to approximately the same MSY value.

11 REFERENCE CASE

11.1 REFERENCE CASE INPUTS

For the reference case runs, all inputs, assumptions, and settings were formulated based on the best available information and scientific judgment. The key settings and any changes from the 2009 reference runs are presented in Table 10. It is important to note that all model runs assumed that Bocaccio productivity, as well as catchability and availability in the surveys and fisheries, were treated as constant over time. While this is not a good assumption for time series that extend over 60 years in some cases (i.e., the halibut fishery effort series), there is no additional information with which to hypothesize credible assumptions on how these parameters varied over time. Where possible, however, we have conducted sensitivity tests to explore the impact of the assumption of stationarity.

11.2 REFERENCE CASE RESULTS

As in the previous work (DFO 2009a)⁶, the results of the reference case indicates that Bocaccio exploitable stock biomass has declined significantly from the 1930s, with the steepest decline occurring from 1985 to 1995 (Table 11, Table 12, Table 13, and Figure 17). The rate of decline slowed after 1995, coincident with lower catches of the early 1990s. The decline appears to have continued after 2000 (Figure 17c).

The posterior mean and median estimates for exploitable biomass in 2012 are 2,205 t and 1,879 t (CV=55%), respectively (Table 11). The posterior median estimate of stock size relative to its unfished stock size (B_{2012}/K) is 3.5% (CV=84%). Current abundance relative to B_{msy} (B_{2012}/B_{msy}) is 7% (CV=84%). The 80% confidence limits (10% and 90% percentiles) of the

⁶ Note that methodology and input for the previous assessment are provided in Stanley et al. 2009a, while final results of the corrected reference run are provided in DFO 2009a.

median estimate of B_{2012}/B_{msy} lie between 0.029 and 0.182 leaving little likelihood that the stock is above the lower PA reference point of $0.4*B_{msy}$ (Figure 18).

The posterior median of F_{2012}/F_{msy} is 1.9 (CV=91%). The maximum rate of increase, r , in the Schaefer model is equal to $2*F_{msy}$, therefore, the population can not sustain values for F_{2012}/F_{msy} of greater than 2. The posterior median for the replacement yield ($RepY$) in 2012 is 143 tons (CV=55%). The posterior median ratio of the total harvest in 2012 relative to replacement yield ($Catch/RepY$) is 99% (CV=87%). Values for this ratio greater than 1 should lead to further stock decline, if sustained.

The reference case median estimates of 2012 biomass and replacement yield are about 80% and 72%, respectively, of those indicated for 2008 (DFO 2009a), with the current of these estimates being more precise (Table 14). The level of depletion (to 2012) is greater than that reported for 2008 (DFO 2009), with the median estimate of stock biomass down to 3.5% of the average unfished level as opposed to 5.6% reported earlier (DFO 2009a). The posterior median ratio of stock size in 2011 relative to 2001 is 0.66 with 10th and 90th percentiles at 0.46 and 0.97 (Table 11). The 90% probability interval for this statistic was 0.42 and 1.09. These results thus indicate that there is more than a 90% probability that stock size is lower in 2011 than it was 10 years previous. The more pessimistic estimates of the current assessment result from a number of reasons. First, the updated priors for survey q were slightly larger owing to the update from the gillnet experiment. The higher and more precise priors for q translate into lower and more precise biomass estimates.

Second, in the 2009 assessment, the survey index values of zero were ignored due to the use of a lognormal likelihood function in that assessment. In the current assessment, we applied a different likelihood function (normal with a constant standard deviation for each index of abundance) that allowed evaluation of zero biomass values. Predictably, the inclusion of zero biomass values for 2000 and, 2011 in the WCVI shrimp survey and for 2007, 2008, and 2010 in the QCSd shrimp survey gives more weight to smaller stock sizes as well as increasing precision.

Error distributions for trawl survey catch rate data are often considered to be more closely approximated by a lognormal distribution than by a normal distribution. In this assessment, we were forced to reject the lognormal density function since it cannot accommodate zero values for observations and there were more than a few of them (five in total). A normal distribution truncated at zero still provided an adequate fit to the abundance indices, as the model converged consistently under a non-linear minimization function. While there were some large positive outliers for some of the time series, the values obtained and applied for the SDs were large enough to accommodate the wide scatter in the deviations between observed and model predicted indices of abundance (Figure 17). In the 2009 assessment, when there were only two zero survey values that were excluded, runs with lognormal versus normal likelihood functions yielded very similar posterior distributions for parameters and stock status.

Table 10. Key parameter choices for the current reference case and significant changes from the 2009 reference case (DFO 2009a, Stanley et al. 2009a).

| Parameter | Value (2012) | Value (2009) | Comments |
|--|---|--|--|
| Prior mean r | 0.1067, SD= 0.039; | 0.117; SD=0.035 | Discussed above |
| Survey values of 0 | Included 0 values | Excluded one 0 value | The 2009 assessment applied a lognormal likelihood function for the abundance index data with a constant standard deviation in the deviation between the logarithms of observed and model predicted abundance index values. For 2012, to accommodate the zero values, we applied a normal distribution for the likelihood function with a constant standard deviation (SD) for each time series of abundance. |
| Recreational catch | Included | Excluded | Described above |
| CVs for indices | Same as 2009 | | As in the 2009 assessment, we applied iterative re-weighting to arrive at values for the standard deviations in deviations between model-predicted and observed abundance index values. |
| Likelihood function for catch | CV=0.6 for troll CV=0.5 for halibut CV=0.4 for recreational | Same for troll and halibut | We applied a truncated normal distribution as the likelihood function for the observed halibut, recreational, and salmon troll catches. We applied a constant fixed CV for each likelihood function. The CVs for the halibut and salmon troll catch values were the same as those applied in the 2009 assessment. The CV for the recreational catch was the value that lead to the closest normal distribution approximation of the expert derived prior for the recreational catch of Bocaccio in 2010. |
| Schaefer surplus production function | $(B_{msy}/K=0.5)$; | Same | Discussed above |
| Salmon troll daily catch | 10 | 15 | Maximum fleet-wide limit on average daily troll catch set at 10 Bocaccio per day. |
| Process error SD | 0.1 | same | |
| Prior mean B_{1935}/K | 0.9 | same | |
| Informative priors for survey q | Updated | same | |
| Density in trawlable area < untrawlable area | Triangular prior updated with experimental data with a median of 1.4. | Triangular prior with mode of 3 but no data used to update it. | Discussed above |
| Process error deviates | Lag = 1 Autocorrelation coeff. = 0.7 | Autocorrelation coeff. = 0.67 | Estimated from posterior median process error deviates, starts in 2012. |

Third, most of the surveys and, particularly the ones that fit the model best, indicate decreases since 2008, despite total catches among the lowest in the history of the fishery. Fourth, exploitation from a fourth sector, the recreational sector, was added in this assessment. As recreational effort has not changed substantially from the 1980s, this fishery is estimated to have exerted a low fishing mortality rate since the 1980s that was not previously included. The addition of this previously unaccounted source of fishing mortality, although relatively small, acts to intensify the estimated decline in stock size.

Table 11. Reference case 2012 stock assessment statistics. Biomass values are in metric tons and the referenced current year is 2012.

| Variable | Mean | SD | CV | 10th Percentile | Median | 90th Percentile |
|----------------------------|--------|--------|-------|-----------------|--------|-----------------|
| r | 0.084 | 0.033 | 0.391 | 0.0397 | 0.084 | 0.1254 |
| B_0 | 63,240 | 38,639 | 0.611 | 26,461 | 52,330 | 116,664 |
| MSY | 1,234 | 904 | 0.733 | 540 | 981 | 2,227 |
| B_{msy} | 31,620 | 19,319 | 0.611 | 13,231 | 26,165 | 58,332 |
| B_{msy}/B_0 | 0.5 | . | . | 0.5 | 0.5 | 0.5 |
| B_{init} | 55,922 | 36,070 | 0.645 | 21,907 | 43752 | 98,206 |
| B_{2012} | 2,205 | 1,214 | 0.55 | 1,031 | 1,879 | 3,625 |
| B_{2012}/B_{msy} | 0.093 | 0.078 | 0.835 | 0.029 | 0.07 | 0.182 |
| B_{2012}/B_{init} | 0.054 | 0.048 | 0.885 | 0.016 | 0.041 | 0.106 |
| B_{2012}/K | 0.047 | 0.039 | 0.835 | 0.0144 | 0.0351 | 0.0911 |
| F_{msy} | 0.0422 | 0.0165 | 0.391 | 0.0199 | 0.042 | 0.0627 |
| F_{2012} | 0.0808 | 0.0359 | 0.444 | 0.041 | 0.0742 | 0.1289 |
| F_{2012}/F_{msy} | 2.2835 | 2.0839 | 0.913 | 1.03 | 1.9037 | 3.5758 |
| $RepY$ | 163 | 90 | 0.552 | 75 | 143 | 287 |
| $Catch/RepY$ | 1.1806 | 1.0267 | 0.87 | 0.5705 | 0.9898 | 1.811 |
| B_{2011}/B_{2001} | 0.6989 | 0.2091 | 0.299 | 0.46 | 0.66 | 0.97 |
| $P(B_{2012} > 0.4B_{msy})$ | 0.01 | | | | | |
| $P(B_{2012} > 0.8B_{msy})$ | 0.001 | | | | | |

Table 12. Posterior 10th, 50th, and 90th percentiles of stock biomass (t) 1935-2012 from the reference case run.

| Year | Lower 10% | Median | Upper 90% | Year | Lower 10% | Median | Upper 90% |
|------|-----------|--------|-----------|------|-----------|--------|-----------|
| 1935 | 21907 | 43752 | 98206 | 1974 | 9813 | 14442 | 24362 |
| 1936 | 21226 | 41662 | 89542 | 1975 | 9223 | 13394 | 23080 |
| 1937 | 20514 | 40226 | 86141 | 1976 | 8556 | 12812 | 22068 |
| 1938 | 19873 | 38530 | 81513 | 1977 | 8514 | 12939 | 21815 |
| 1939 | 19151 | 37378 | 80921 | 1978 | 8816 | 12815 | 21613 |
| 1940 | 18697 | 34925 | 75919 | 1979 | 8586 | 12868 | 21257 |
| 1941 | 18386 | 33731 | 66959 | 1980 | 8369 | 13031 | 21236 |
| 1942 | 17717 | 33074 | 68540 | 1981 | 8650 | 12822 | 20570 |
| 1943 | 17250 | 31801 | 61479 | 1982 | 8810 | 12708 | 20311 |
| 1944 | 16862 | 31125 | 59523 | 1983 | 8695 | 12730 | 19853 |
| 1945 | 16925 | 30524 | 58881 | 1984 | 8489 | 12407 | 18751 |
| 1946 | 16222 | 29843 | 57254 | 1985 | 8512 | 11886 | 17679 |
| 1947 | 15891 | 28407 | 54899 | 1986 | 8278 | 11095 | 16151 |
| 1948 | 15565 | 27689 | 53313 | 1987 | 7540 | 10351 | 14590 |
| 1949 | 15046 | 26894 | 52294 | 1988 | 6706 | 9123 | 12871 |
| 1950 | 14438 | 26449 | 51442 | 1989 | 5957 | 8023 | 11385 |
| 1951 | 13948 | 25297 | 49527 | 1990 | 5275 | 7153 | 10404 |
| 1952 | 13618 | 24702 | 47783 | 1991 | 4672 | 6202 | 9207 |
| 1953 | 13809 | 24445 | 45886 | 1992 | 3914 | 5327 | 7900 |
| 1954 | 13992 | 24667 | 44708 | 1993 | 3090 | 4401 | 6705 |
| 1955 | 14039 | 24184 | 44839 | 1994 | 2498 | 3638 | 5773 |
| 1956 | 13587 | 23946 | 44413 | 1995 | 2149 | 3240 | 5203 |
| 1957 | 13314 | 23410 | 43870 | 1996 | 1917 | 2930 | 4864 |
| 1958 | 13290 | 23189 | 41641 | 1997 | 1843 | 2829 | 4810 |
| 1959 | 12828 | 22328 | 41258 | 1998 | 1843 | 2709 | 4650 |
| 1960 | 12873 | 22438 | 40503 | 1999 | 1832 | 2728 | 4537 |
| 1961 | 13014 | 21777 | 39103 | 2000 | 1842 | 2749 | 4526 |
| 1962 | 13193 | 21277 | 38402 | 2001 | 1825 | 2718 | 4596 |
| 1963 | 12938 | 21363 | 37243 | 2002 | 1726 | 2599 | 4484 |
| 1964 | 12805 | 21415 | 36138 | 2003 | 1609 | 2502 | 4253 |
| 1965 | 13015 | 20672 | 35076 | 2004 | 1558 | 2475 | 4236 |
| 1966 | 12561 | 20286 | 33724 | 2005 | 1458 | 2428 | 4233 |
| 1967 | 12024 | 18819 | 32355 | 2006 | 1443 | 2365 | 4157 |
| 1968 | 11845 | 18224 | 31859 | 2007 | 1339 | 2278 | 4030 |
| 1969 | 11307 | 17697 | 30050 | 2008 | 1270 | 2230 | 3941 |
| 1970 | 11103 | 16759 | 28693 | 2009 | 1178 | 2071 | 3676 |
| 1971 | 11026 | 16194 | 27682 | 2010 | 1082 | 1935 | 3435 |
| 1972 | 10971 | 16118 | 26671 | 2011 | 1052 | 1911 | 3506 |
| 1973 | 10463 | 15279 | 26040 | 2012 | 1031 | 1879 | 3625 |

Table 13. Posterior 10th, 50th, and 90th percentiles of the ratio of stock biomass in each recent year to the stock biomass sixty years prior.

| Year | 10% | 50% | 90% |
|------|------|------|------|
| 1994 | 0.03 | 0.07 | 0.15 |
| 1995 | 0.03 | 0.07 | 0.15 |
| 1996 | 0.03 | 0.07 | 0.15 |
| 1997 | 0.04 | 0.07 | 0.15 |
| 1998 | 0.04 | 0.07 | 0.15 |
| 1999 | 0.04 | 0.08 | 0.16 |
| 2000 | 0.04 | 0.08 | 0.16 |
| 2001 | 0.04 | 0.08 | 0.16 |
| 2002 | 0.04 | 0.08 | 0.16 |
| 2003 | 0.04 | 0.08 | 0.16 |
| 2004 | 0.04 | 0.08 | 0.17 |
| 2005 | 0.04 | 0.07 | 0.16 |
| 2006 | 0.04 | 0.07 | 0.16 |
| 2007 | 0.04 | 0.07 | 0.15 |
| 2008 | 0.04 | 0.07 | 0.15 |
| 2009 | 0.03 | 0.07 | 0.16 |
| 2010 | 0.03 | 0.07 | 0.16 |
| 2011 | 0.03 | 0.07 | 0.15 |
| 2012 | 0.03 | 0.07 | 0.15 |

Table 14. Comparison of key results from the 2009 (DFO 2009a) and 2012 analyses. B_{cur} refers either to 2012 or 2008.

| Variable | 2012 | | | 2009 | | |
|-------------------|--------|-------|--------|--------|---------|--------|
| | Mean | CV | Median | Mean | Mean CV | Median |
| B_0 | 63,240 | 0.611 | 52,330 | 54,042 | 0.66 | 45,053 |
| B_{msy} | 31,620 | 0.611 | 26,165 | 27,021 | 0.662 | 22,526 |
| B_{cur} | 2,205 | 0.55 | 1,879 | 3,022 | 0.83 | 2,324 |
| B_{cur}/B_{msy} | 0.093 | 0.835 | 0.070 | 0.155 | 0.973 | 0.111 |
| B_{cur}/K | 0.047 | 0.835 | 0.0351 | 0.078 | 0.973 | 0.056 |
| $RepY$ | 163 | 0.552 | 143 | 236 | 0.649 | 198 |




Figure 17. Plots of median and 80% probability intervals and indices rescaled by their median a. 1935-2012; b. 1975-2012; c. 2000-2012. Note that some of the very large values for some of the indices are not shown in panels b. and c. to permit closer inspection of more recent trends.




Figure 18. Estimated stock status in 2012 and exploitation rate relative to reference points (from Table 11). Range indicates 10th and 90th percentiles.

11.3 REFERENCE CASE MODEL PERFORMANCE AND UNCERTAINTY

Posterior distributions for most quantities show an update from the prior distributions (Figure 19). The post model, pre-data distributions were also shown in Figure 19. The post model, pre-data distribution shows how the priors interact with the BSP model, fixed inputs for catch, and fishing effort for the different imputed fisheries before the model is fitted to the abundance index data. The post model, pre-data distributions indicate that the priors for model parameters, when applied in combination with the inputted values for catch and effort, provide quite vague information about most of the model parameters and quantities of interest. While the post model, pre-data distribution for catch to replacement yield in 2012 appears to be informed by the model inputs and model structures, the range is still quite wide with most outputted values ranging between about 0.1 and 2 and is not updated after fitting to the abundance indices (Figure 19). This is the only posterior distribution that is not significantly updated by the abundance data.

Model fits to the survey and CPUE data are poor, with large deviations between observed and predicted indices. CVs for the predicted to observed fits to the abundance indices are greater than 0.5 for seven out of nine series (Table 15; Figure 17). This outcome is caused by some large outlier values in both shrimp trawl series and in the US Triennial series (Figure 12). Autocorrelation is apparent in the deviates for some of the indices, as, for example, in the US Triennial series (Figure 17). The posterior mean for the intrinsic rate of increase (r) was 0.084 (CV=39%), less than the prior of 0.107 (CV=37%) (Figure 19c). This decrease in the mean value for r suggests that the model reduced the average underlying stock productivity in order to fit the recent declines in biomass.

It is tempting to configure the model to fit the US Triennial index more closely because it indicates an intuitively acceptable monotonic trend. However, this is not only a circular argument but, as noted in Stanley et al. (2009a), the apparent trend is highly leveraged by one

anomalous tow in the first year. We suggest that the reference case weighting of all indices is appropriate in not allowing single survey points (or surveys) to have undue influence.

The marginal posterior distributions indicated that moderate amounts of precision were obtained for most parameters (Figure 19). However, the large skews and long tails remain for some estimates. For example, much of the probability for carrying capacity lies well below 75,000 t while the tail stretches to 175,000 tonnes. Estimates for some other quantities are well defined. For example, for B_{2012}/B_0 , the majority of the probability lies below 10% of B_0 .

The annual process error deviates from the predicted surplus production were strongly negative for 2006-2009 (Figure 20). This indicates that the model production function predicted higher production in these years than has been reflected in the surveys. This effect is in addition to the lowering of the average stock productivity discussed in the previous paragraph, indicating that recent stock productivity has been even lower than predicted, even for an average $r=0.084$. Although none of the deviates was significantly different from 0, these negative deviates in 2006-2009 suggest that there was poor recruitment into exploitable age classes in these years. This relatively poor recruitment may explain, in part, why the population has not responded to the recent reduction in catches.

For the reference case and other model runs, the autocorrelation coefficient at lag 1 in the process error deviates from 1980 to 2010 was estimated at about 0.7, which was significant ($p<0.05$). This implies that there is a strong tendency for a poor year of surplus production to be followed by poor years.




Figure 19. Marginal posterior distributions for a. carrying capacity (K or B_0); b. maximum sustainable yield (m_{sy}); c. r ; d. replacement yield in 2012; e. stock biomass in 2012; f. the ratio of fishing mortality rate in 2012 to that at m_{sy} ; g. the ratio of stock biomass in 2012 to average unfished stock size; and, h. the ratio of total catch biomass in 2012 to replacement yield in 2012. Priors are shown for B_0 and r . The post model, pre-data distributions are shown for derived quantities to show the influence of the catch and effort data, model structure, and prior distributions on model output distributions for quantities of interest.

Table 15. Values for the CVs applied for each of the abundance indices in the reference case and other model runs. Note that the CVs derived from iterative reweighting changed little between runs.

| Index | Number of data points | Years | Standard deviation | Average value | Approx. CV |
|-------------------|-----------------------|-----------|--------------------|---------------|------------|
| WCVI GF | 4 | 2004-2010 | 135 | 229 | 0.59 |
| QCSd GF | 6 | 2003-2011 | 110 | 161 | 0.68 |
| HS GF | 4 | 2005-2011 | 25 | 35 | 0.71 |
| WCHG GF | 4 | 2006-2010 | 2 | 10 | 0.20 |
| WCVI shrimp trawl | 35 | 1975-2011 | 170 | 123 | 1.38 |
| QCSd shrimp trawl | 13 | 1999-2011 | 135 | 43 | 3.14 |
| US Triennial | 7 | 1981-2001 | 1,900 | 2,176 | 0.87 |
| Comm. CPUE | 8 | 1996-2003 | 4.5 | 29 | 0.15 |
| IPHC | 9 | 2003-2011 | 1.1 | 2 | 0.51 |




Figure 20. Annual deviates in surplus production with the median and 80% probability intervals shown. Estimates are shown only for years after 1975 because without data prior to then, the posteriors for these deviates are determined by the prior.

11.4 REFERENCE CASE RECONSTRUCTION OF HISTORICAL CATCHES

The estimated historical catch for the halibut, salmon troll, and recreational fisheries from the reference case are provided in Figure 21, with the median values and the estimated relative contributions from each sector provided in Appendix Table 1 and Appendix Table 2.

The posterior distributions for the catchability coefficients for the halibut, salmon troll, and recreational fisheries show some moderate updating from the priors to favour smaller values (Figure 22). The posterior distributions still show considerable uncertainty with long right hand tails, especially for the halibut and recreational fisheries. The more precise distribution for the salmon troll fishery results from the extremely high historic effort relative to current effort and the estimated high catches for this fishery prior to the mid-1990s.




Figure 21. Median and 80% probability intervals for catch of Bocaccio in the: a. halibut fishery, b. salmon troll fishery, and c. recreational fishery, d. all sectors combined including trawl and ZN HL. Note the large variation in scale on the y-axis.




Figure 22. Prior and posterior density functions for the catchability coefficients for the a. halibut, b. salmon troll, and c. recreational sectors.

12 SENSITIVITY TESTS

12.1 SENSITIVITY RUNS

Eighteen additional runs were used to evaluate the sensitivity of the results in the reference case to alternative assumptions (Table 16) (see also Stanley et al. 2009a in which 31 additional model runs were examined).

A common point of uncertainty with the implementation of the Schaefer model is its assumption that B_{msy}/B_0 rests at 0.5. While this may be plausible for a population with very low productivity (low maximum intrinsic rate of increase), alternative values cannot be ruled out. We therefore examined runs with B_{msy}/B_0 fixed at values of 0.3, 0.4, and 0.6 (**A.1-A.3**).

The reference prior for r , was formed using empirically derived uncertainty distributions for all life history parameters; however, there is uncertainty in these inputs. One of these includes the posterior predictive distribution for the Ricker steepness parameter for rockfishes (Forrest et al. 2010). In DFO 2009a, two cases examined prior medians for r at 67% and 133% of the

reference case prior. With a higher input distribution for steepness, the prior CV in r remained the same and the prior standard deviation increased proportionally with the mean value for r .

We captured the uncertainty in r by specifying two alternative prior distributions for r by setting the prior mean to 67% of the reference case prior mean (low prior r) and setting the prior mean to 133% of the reference case prior mean (high prior r). The prior CV was held constant for both cases (**B.1 and B.2**).

Table 16. Summary of sensitivity test runs.

| Code | Category Description | Code | Run Description |
|------|----------------------------------|------|--|
| Ref | Reference run | Ref | Reference run |
| A | B_{msy}/K | A.1 | $B_{msy}/K = 0.3$ |
| | | A.2 | $B_{msy}/K = 0.4$ |
| | | A.3 | $B_{msy}/K = 0.6$ |
| B | r prior mean | B.1 | low r (mean = 0.0802, SD = 0.0391) |
| | | B.2 | High r (mean = 0.142, SD = 0.052) |
| C | Catch assumptions | C.1 | Sum of trawl and non-halibut hook and line catch x 0.5 all yrs |
| | | C.2 | Sum of trawl and non-halibut hook and line catch x 1.5 all yrs |
| | | C.3 | Sum of trawl and non-halibut hook and line catch x 0.25 86-95 |
| | | C.4 | Sum of trawl and non-halibut hook and line catch x 0.5 86-95 |
| D | Survey q priors | D.1 | Non-informative priors for all constants of proportionality for abundance indices (q); priors for the catchability coefficients for the imputed fisheries were kept the same as in the reference case, i.e., also non-informative. |
| E | Effect of data | E.1 | Include only one data point per series and non-informative priors for all q s: post model, pre-data analysis of output distributions. |
| F | Bycatch assumptions ¹ | F.1 | Halibut catchability for Bocaccio decreased by 2.0% per year (implies effort in 1935 4.7 x reference effort in 1935) |
| | | F.2 | Halibut catchability for Bocaccio decreased by 1.5% per year (implies effort in 1935 3.2 x reference effort in 1935) |
| | | F.3 | Halibut catchability for Bocaccio decreased by 1% per year (implies effort in 1935 2.2 x reference effort in 1935) |
| | | F.4 | Halibut catchability for Bocaccio increased by 1% per year (implies effort in 1935 0.5 x reference effort in 1935) |
| | | F.5 | Halibut catchability for Bocaccio increased by 1.5% per year (implies effort in 1935 0.3 x reference effort in 1935) |
| | | F.6 | Halibut catchability for Bocaccio increased by 2.0% per year (implies effort in 1935 0.2 x reference effort in 1935) |
| | | F.7 | Model started in 1900 with 1900–1934 halibut effort assumed proportional to halibut catch in same year, scaled to the 1935 halibut catch and effort; other catch and effort series set=0 in 1900 and increased proportionately to 1935 observed values |

¹ Sensitivity runs F.1 to F.6 were generated relative to average absolute effort observed from 2006 to 2011.

The uncertainty in the fixed catch estimates for trawl and HL fisheries was captured by investigating two alternative scenarios which set these historic catches to 50% and 150% of the reference case, respectively (**C.1 and C.2**). Two additional sensitivity runs were performed in which the fixed catch estimates for domestic trawling in 1986-1995 were decreased by a factor of 0.25 and 0.5. This was to reflect the possibility that sales slip and fisher logbook data inflated the catches of Bocaccio in this period (**C.3 and C.4**). The sensitivity of the model results to informed priors on q was investigated by replacing these priors with uninformative priors with wide bounds (uniform over $\ln(q)$) (**D.1**).

A question often arises in Bayesian stock assessment about the degree to which the model structure, acting with priors and the fixed inputs, influences the stock status results. To address

this issue, we evaluated the influence of the reference case priors and fixed values for trawl and HL catch and fishing effort. We did this by producing what has been called the "post-model pre-data distribution" of model outputs (Punt and Butterworth 2002). We ran the model by informing it with the prior distributions for all of the estimated parameters and applying the fixed input values for catch and historic troll, halibut, and recreational effort but without fitting the model to the abundance data (E.1). In other words, we effectively drew values for parameters from their prior distributions and projected the model with the fixed catch and effort values. We then compiled the frequency distribution of outputted parameter values and quantities of interest. We did so without weighting the trajectories according to how well the modelled trajectories fitted the abundance indices and the data on catch for the halibut, salmon troll, and recreational fisheries. Even without the likelihood function applied, we can expect some updates to the prior distributions. Some combinations of parameter values drawn from the prior distributions will result in population trajectories that crash the population before the current year. These runs are weeded out and not counted in tallying up the post-model, pre-data distributions.

A strong assumption in the imputation of bycatch based on a time series of effort is that the catchability (k) of Bocaccio in these fisheries has remained constant over time. The fishery with the largest imputed bycatch of Bocaccio is the halibut fishery therefore; we carried out a number of sensitivity runs based on assumed constant rates of change in k over the time series. We carried out six additional model runs where k was modelled to change at rates of -2%, -1.5%, -1%, 1%, 1.5%, and 2% per year (F.1-F.6). In addition, the halibut effort in the initial year, 1935, starts out high and gradually drops. Early records of halibut catches in BC (Bell et al. 1952) show that catches were very low in 1900, followed by a gradually increasing trend, peaking in the 1920s and subsequently dropping by about 40% in the 1930s (Table 17). We carried out an additional sensitivity run in which the model was started in 1900. This run imputed the values for halibut effort based on the assumption that halibut effort was directly proportional to halibut catch during the period 1900 to 1934, scaled relative to the 1935 halibut catch and effort. Other catch and effort series were filled by starting each series at zero and increasing each proportionately to reach the observed values in 1935 (F.7).

The prior mean for P_0 (i.e., the ratio of stock biomass in the initial year to B_0) was set at 0.9, with a prior coefficient of variation (CV) of about 0.2, as in the 2009 assessment. While the actual uncertainty in P_0 may be greater, numerous studies have shown that, providing the stock assessment model starts several decades in the past as this one does, the prior mean presumed for P_0 , has very little effect on estimates of key parameters and stock status (e.g., Stanley et al. 2009a; King et al. 2012; Yamanaka et al. 2012). Due to this, there was no need to include different priors for P_0 in the sensitivity analyses.

No sensitivity runs were made to explore possible changes in species productivity over the model period. It is certainly plausible that average Bocaccio productivity has varied over the 60-70 years that are modelled in this analysis; however, we have no specific information to assist us in modelling time-dependent changes in the appropriate parameters. Furthermore, only if future variation in productivity was predictable, would such modelling assist in managing the Bocaccio population.

Table 17. Records of Pacific Halibut catch (t) in BC waters from 1900-1937 (Bell et al. 1952).

| Year | Catch | Year | Catch |
|------|--------|------|--------|
| 1900 | 3,598 | 1919 | 20,084 |
| 1901 | 4,998 | 1920 | 23,233 |
| 1902 | 7,312 | 1921 | 29,892 |
| 1903 | 9,062 | 1922 | 26,906 |
| 1904 | 12,180 | 1923 | 30,029 |
| 1905 | 7,200 | 1924 | 29,997 |
| 1906 | 9,950 | 1925 | 29,547 |
| 1907 | 12,915 | 1926 | 27,681 |
| 1908 | 15,892 | 1927 | 26,786 |
| 1909 | 19,460 | 1928 | 30,467 |
| 1910 | 19,387 | 1929 | 28,656 |
| 1911 | 15,854 | 1930 | 24,466 |
| 1912 | 21,127 | 1931 | 18,374 |
| 1913 | 22,347 | 1932 | 17,046 |
| 1914 | 21,444 | 1933 | 17,027 |
| 1915 | 31,769 | 1934 | 18,313 |
| 1916 | 26,723 | 1935 | 17,129 |
| 1917 | 23,030 | 1936 | 17,001 |
| 1918 | 17,793 | 1937 | 18,917 |

12.2 SENSITIVITY RESULTS

In general, the sensitivity tests did not reveal any significantly different stock status conclusions relative to the reference case (Table 18). All of the runs, except for the diagnostic run, **E.1**, continue to indicate a stock that is well below $0.4*B_{msy}$ and that current catch levels are approximately equal to estimates of replacement yield (*RepY*).

Results were relatively insensitive to the choice of B_{msy}/K (**A.1-A.3**). Posterior median values for B_{curr}/B_{msy} increased from 0.076 to 0.110 when the B_{msy}/K ratio decreased from 0.6 to 0.3, which was the greatest range observed in stock status among the 18 sensitivities investigated. The estimates of the ratios of total catch to replacement yield (*Catch/RepY*), and the current fishing mortality rate to F_{msy} (F_{curr}/F_{msy}) were also relatively insensitive to choice of B_{msy}/K , although these estimates were the least optimistic for the highest setting of B_{msy}/K , as expected.

The posterior median value for B_{msy} was largest for the reference case with B_{msy}/K set at 0.5. In contrast, one might think it should be largest for the run with the largest B_{msy}/K (i.e., 0.6). The smaller estimate of B_{msy} for the 0.6 run results mainly from a marked discontinuity in the shape of the Fletcher generalized production function as B_{msy}/K increases from below to above 0.5 as discussed earlier (Figure 16). The data to which the surplus production models were fitted tend to reference the different production curves to the similar *msy* values, since we see little change in abundance for several years at similar levels of low catches after strong depletion. At the same value for *msy*, the Schaefer model with B_{msy}/K at 0.5 predicts the largest values for B_{msy} and K when compared to the Fletcher model variants with different B_{msy}/K . Below the value of 0.5 for B_{msy}/K , B_{msy} decreases with B_{msy}/K . Similarly, values of B_{msy}/K above 0.5 lead to decreases in B_{msy} and K . This is a mathematical consequence of the Fletcher model parameterization (Quinn and Deriso 1999).

Overall results varied modestly and predictably with different priors for r (**B.1 and B.2.**), and high and low scenarios catch scenarios (**C.1 and C.2**). For example, lower values for r and higher values for catches scaled up the estimates of current stock biomass but the estimates of depletion and *Catch/RepY* in 2012 remained insensitive to these alternative input settings. The

lower catch scenario slightly reduced the estimate of *Catch/RepY*. When the fixed catch inputs for 1986-1995 were lowered by factors of four and two (C.3 and C.4), the effect was to slightly reduce the posterior mean for r , reduce the estimated replacement yield in recent years, and lower the estimated current stock status for run C.3. It remained the same as the reference case for run C.4 (Table 18).

Application of a non-informative prior for q resulted in considerably lower precision in the estimates of biomass-related quantities (**D.1**). For example, the 80% probability range for biomass in 2012 was 970-5,500t versus 1,000-3,600t under the non-informative and informative priors, respectively. The posterior median values with the non-informative prior were slightly less pessimistic mainly because of the large positive skew for biomass under the non-informative prior for q . For example, *Catch/RepY* was 0.83 compared to 0.99 under the reference case.

The removal of all of the stock assessment data under the reference case to create a post model, pre-data run (**E.1**), yielded much wider probability distributions for all biomass derived quantities and show the influence of the priors on model output distributions when the fixed values for historic catch and effort are applied. For example, the 80% probability interval for stock biomass in 2012 widened to 3,600-81,000 t as compared with 1,000-3,600 t under the reference case. The confidence interval for B_{2012}/B_{msy} increased to 0.07-1.4 from 0.03-0.18 under the reference case. This run shows that the model structure, priors, and fixed inputs for catch and effort acting together, do not lend high precision to any of the results. Nor do they strongly bias the stock status results in one direction. The precision in the estimates of status derive mainly when the model is fitted to the abundance index data and data on catch for the three fisheries with catches estimated from effort series.

Estimated stock status showed a slight increasing trend from F.1 (catchability decreasing by 2%/year) to F.6 (catchability increasing by 2%/year) (Table 18). However, the magnitude of this increase was slight, demonstrating that the model conclusions regarding stock status were insensitive to the assumptions for the annual rate of change in the catchability coefficient, k . The magnitude of the estimated Bocaccio catch in the early years of the fishery varied considerably as a result of the different assumptions used for k and the consequent variation in the time series of effective halibut effort (Figure 23). Extending the model backwards to 1900 resulted in posterior distributions for all parameters that were nearly indistinguishable from the reference case. This implied little sensitivity to the choice of beginning year for the stock reconstruction with respect to estimates of current biomass level and relative stock status.

To provide a better understanding on how the different sensitivity runs behaved, the posterior medians for B_{init}/K , ($init=1935$), the posterior median estimated catches for the halibut, troll, and recreational fisheries, total catch, replacement yield and the posterior median for the total catch to replacement yield are also provided (Table 19). The posterior median values for the B_{init}/K ratio values were largely determined by the prior distribution and were similar across all model runs.

The posterior medians for the estimated catches were also similar across runs, except for the post model, pre-data run (**E.1**) which gave very high values for output catch and stock biomass distributions. This occurred because the reference catch data for the imputed fisheries were ignored and not used to weight the different model trajectories in the computation of model output distributions. As mentioned above, the absolute level of imputed bycatch for the halibut fishery from 1935 was sensitive to the values assumed for the catchability parameter k .

The posterior median for total catch was less than the posterior median for replacement yield for the instances in which the Fletcher-Schaefer model was run. In **A.1**, while the posterior median of *Catch/RepY* was about 1, the posterior median for the total catch (148 t) was less than the posterior median for the replacement yield (176 t) (Table 19, Figure 24). This was a

consequence of the strong discontinuity in the shape of the Fletcher-Schaefer production function that was applied in this run (Figure 16). The same pattern and explanation applies to **A.2**. The resulting differences in the posterior distributions for *RepY* and total catch, and somewhat ragged relationship between total catch and *RepY* in these runs are illustrated in Figure 24. Note that median estimates of ratio *Catch/RepY* will not necessarily equal the ratio of median estimates of catch and median estimate of *RepY*.

Table 18. Medians and 80% credibility intervals drawn from the posterior distributions for seven parameters taken from the Bocaccio assessment for the reference run and 18 sensitivity runs. Codes used for each run along with a run description can be found in Table 16. Biomass values are in tons.

| Run | <i>r</i> | | | <i>B_{msy}</i> | | | <i>B_{current}</i> | | | <i>RepY</i> | | | <i>B_{current}/B_{msy}</i> | | | <i>F_{current}/F_{msy}</i> | | | <i>Catch_{curr}/RepY</i> | | |
|------|---------------------------------|--------|-------|------------------------|--------|-------|----------------------------|--------|-------|-------------|--------|------|--|--------|------|--|--------|------|----------------------------------|--------|------|
| | 10% | Median | 90% | 10% | Median | 90% | 10% | Median | 90% | 10% | Median | 90% | 10% | Median | 90% | 10% | Median | 90% | 10% | Median | 90% |
| Ref. | 0.040 | 0.084 | 0.125 | 13231 | 26165 | 58332 | 1031 | 1879 | 3625 | 75 | 143 | 287 | 0.029 | 0.070 | 0.18 | 1.03 | 1.90 | 3.58 | 0.57 | 0.99 | 1.81 |
| | <i>B_{msy}/K</i> | | | | | | | | | | | | | | | | | | | | |
| A.1 | 0.052 | 0.097 | 0.152 | 9677 | 19596 | 37601 | 1072 | 2077 | 3917 | 81 | 176 | 409 | 0.048 | 0.110 | 0.28 | 1.02 | 1.89 | 3.53 | 0.56 | 1.00 | 1.84 |
| A.2 | 0.047 | 0.093 | 0.152 | 10665 | 23007 | 46577 | 1071 | 1969 | 3885 | 74 | 170 | 414 | 0.039 | 0.085 | 0.23 | 1.07 | 1.96 | 3.63 | 0.59 | 1.03 | 1.90 |
| A.3 | 0.038 | 0.078 | 0.123 | 11512 | 24340 | 60034 | 916 | 1878 | 3753 | 63 | 140 | 291 | 0.028 | 0.076 | 0.21 | 1.06 | 2.10 | 4.58 | 0.57 | 1.05 | 2.17 |
| | <i>r</i> prior mean | | | | | | | | | | | | | | | | | | | | |
| B.1 | 0.022 | 0.06 | 0.104 | 15192 | 31551 | 65200 | 1228 | 2185 | 4072 | 48 | 122 | 241 | 0.030 | 0.067 | 0.17 | 1.14 | 2.20 | 5.58 | 0.63 | 1.15 | 2.85 |
| B.2 | 0.04 | 0.094 | 0.157 | 9970 | 18703 | 39430 | 907 | 1716 | 3447 | 66 | 146 | 277 | 0.037 | 0.092 | 0.22 | 0.91 | 1.80 | 4.03 | 0.53 | 0.948 | 2.02 |
| | Catch assumptions | | | | | | | | | | | | | | | | | | | | |
| C.1 | 0.033 | 0.074 | 0.115 | 9214 | 17717 | 34973 | 671 | 1205 | 2428 | 40 | 81 | 162 | 0.032 | 0.065 | 0.18 | 0.94 | 1.83 | 3.46 | 0.53 | 0.97 | 1.76 |
| C.2 | 0.049 | 0.089 | 0.135 | 15977 | 30036 | 56272 | 1312 | 2297 | 4510 | 97 | 192 | 363 | 0.032 | 0.073 | 0.19 | 1.04 | 2.012 | 3.99 | 0.59 | 1.04 | 2.01 |
| C.3 | 0.034 | 0.071 | 0.108 | 14406 | 26808 | 51374 | 1030 | 1735 | 3442 | 58 | 119 | 233 | 0.028 | 0.063 | 0.16 | 1.267 | 2.37 | 4.86 | 0.69 | 1.23 | 2.40 |
| C.4 | 0.039 | 0.078 | 0.119 | 12970 | 25326 | 51962 | 1041 | 1897 | 3599 | 63 | 135 | 271 | 0.032 | 0.075 | 0.19 | 1.09 | 2.10 | 4.37 | 0.61 | 1.09 | 2.18 |
| | Survey <i>q</i> priors | | | | | | | | | | | | | | | | | | | | |
| D.1 | 0.051 | 0.094 | 0.144 | 11629 | 20908 | 49517 | 969 | 2208 | 5450 | 82 | 181 | 404 | 0.036 | 0.098 | 0.30 | 0.697 | 1.53 | 3.37 | 0.42 | 0.83 | 1.68 |
| | Effect of data | | | | | | | | | | | | | | | | | | | | |
| E.1 | 0.086 | 0.132 | 0.18 | 28358 | 59382 | 89656 | 3589 | 24012 | 80996 | 393 | 1944 | 4777 | 0.069 | 0.522 | 1.38 | 0.353 | 1.03 | 2.39 | 0.41 | 0.84 | 1.41 |
| | Assumptions about catch | | | | | | | | | | | | | | | | | | | | |
| F.1 | 0.041 | 0.082 | 0.124 | 16897 | 34823 | 69233 | 1227 | 2212 | 4070 | 85 | 164 | 310 | 0.027 | 0.060 | 0.17 | 0.92 | 1.71 | 3.21 | 0.51 | 0.89 | 1.60 |
| F.2 | 0.044 | 0.086 | 0.120 | 16375 | 32473 | 66401 | 1106 | 1997 | 3800 | 85 | 148 | 284 | 0.027 | 0.058 | 0.16 | 1.01 | 1.89 | 3.21 | 0.56 | 0.99 | 1.63 |
| F.3 | 0.044 | 0.082 | 0.124 | 14879 | 29374 | 59814 | 1101 | 1944 | 3754 | 77 | 149 | 280 | 0.030 | 0.064 | 0.18 | 1.00 | 1.89 | 3.55 | 0.56 | 0.98 | 1.78 |
| F.4 | 0.045 | 0.089 | 0.129 | 12261 | 21667 | 50526 | 843 | 1882 | 3519 | 74 | 145 | 293 | 0.034 | 0.074 | 0.20 | 1.05 | 1.96 | 3.73 | 0.58 | 1.03 | 1.89 |
| F.5 | 0.044 | 0.087 | 0.130 | 11529 | 21187 | 48401 | 932 | 1840 | 3642 | 68 | 138 | 282 | 0.034 | 0.078 | 0.20 | 1.07 | 2.08 | 4.07 | 0.60 | 1.09 | 2.03 |
| F.6 | 0.043 | 0.088 | 0.135 | 11477 | 19427 | 40977 | 976 | 1847 | 3592 | 70 | 148 | 282 | 0.040 | 0.089 | 0.21 | 1.04 | 1.94 | 3.90 | 0.59 | 1.03 | 1.96 |
| F.7 | 0.044 | 0.086 | 0.132 | 13226 | 26644 | 60055 | 1000 | 1930 | 3709 | 76 | 144 | 296 | 0.030 | 0.068 | 0.19 | 0.98 | 1.87 | 3.65 | 0.55 | 0.98 | 1.83 |

Table 19. Posterior medians for estimated catch values and replacement yield statistics for 2012 taken from the Bocaccio assessment for the reference run and a selection of the sensitivity runs. Codes used for each run along with a run description can be found in Table 16. Catch values are in tonnes.

| Run | B_{init}/K | Halibut catch (1935) | Halibut catch (2012) | Salmon troll catch (2012) | Rec. catch (2012) | Total catch (2012) | RepY (2012) | Catch/RepY (2012) |
|------|--------------|----------------------|----------------------|---------------------------|-------------------|--------------------|-------------|-------------------|
| Ref. | 0.88 | 1,242 | 6 | 6 | 7 | 140 | 143 | 0.99 |
| A.1 | 0.89 | 1,366 | 6 | 6 | 9 | 148 | 176 | 1.00 |
| A.2 | 0.87 | 1,153 | 5 | 6 | 8 | 145 | 170 | 1.03 |
| A.3 | 0.89 | 981 | 6 | 7 | 7 | 144 | 140 | 1.05 |
| B.1 | 0.88 | 1336 | 6 | 6 | 6 | 140 | 122 | 1.14 |
| B.2 | 0.87 | 584 | 4 | 6 | 6 | 137 | 146 | 0.95 |
| C.1 | 0.88 | 797 | 4 | 7 | 6 | 77 | 81 | 0.97 |
| C.2 | 0.89 | 1,247 | 7 | 6 | 6 | 200 | 192 | 1.04 |
| C.3 | 0.87 | 1,364 | 6 | 9 | 7 | 144 | 119 | 1.23 |
| C.4 | 0.87 | 1,172 | 6 | 8 | 7 | 144 | 135 | 1.09 |
| D.1 | 0.89 | 992 | 6 | 8 | 7 | 146 | 181 | 0.83 |
| E.1 | 0.87 | 2,422 | 127 | 50 | 964 | 1,407 | 1,944 | 0.84 |
| F.1 | 0.88 | 4,872 | 4 | 7 | 7 | 141 | 164 | 0.89 |
| F.2 | 0.90 | 3,639 | 5 | 7 | 8 | 142 | 148 | 0.99 |
| F.3 | 0.89 | 2,422 | 5 | 7 | 7 | 141 | 149 | 0.98 |
| F.4 | 0.87 | 620 | 7 | 6 | 7 | 146 | 145 | 1.03 |
| F.5 | 0.89 | 434 | 7 | 6 | 7 | 146 | 138 | 1.09 |
| F.6 | 0.89 | 294 | 8 | 7 | 7 | 147 | 148 | 1.03 |
| F.7 | 0.87 | 910 | 5 | 5 | 7 | 142 | 144 | 0.98 |




Figure 23. Plots of a. effective halibut effort in BC waters and b. posterior median catch of Bocaccio by halibut gear under different scenarios for constant percent changes in catchability of halibut gear for Bocaccio, (i.e., termed “technological creep” or “TC”).




Figure 24. Plots of the a. marginal posterior distributions for total catch and replacement yield; b. draws from the posterior distribution of values for replacement yield versus total catch from the reference case run; c. marginal posterior distributions for total catch and replacement yield and d. draws from the posterior distributions of values for replacement yield versus total catch from run A.1.

12.3 SENSITIVITY RUNS EVALUATED WITH BAYES FACTORS

To compare the credibility of each model given the data, we computed Bayes factors (Kass and Raftery 1995) for the reference case and for each of the related sensitivity runs. Bayes factors account for both the relative goodness of fit of the model to the data and the parsimony for each of the alternative models. They are calculated as the ratio of the marginal probability of the data for one model to that for another model. We used the mean value for the importance weights from a given model run as an approximation of the probability of the data given the model (Kass and Raftery 1995, McAllister and Kirchner 2002). This is known to be a numerically stable approximation for the probability of the data, given the model and approximations obtained through importance sampling. For example, the CV in the natural logarithm in the mean weight was less than 0.05 after several million draws from the importance function. In all instances, we compared Bayes factors to our reference case model settings. In other words, the probability of the data for the reference case model was placed in the denominator and that for the model run to which it was compared in the numerator. It is commonly held that the Bayes factor must depart substantially from 1.0 for anything to be inferred from the exercise but even fairly large or small departures in Bayes factors can result from random chance in the data and/or misspecification of probability models. Intermediate values for Bayes factor (e.g., between about 0.001 and 100) should be interpreted with caution. For example, models that had Bayes factors of between about 0.1 and 0.01 could be interpreted as unlikely but not discredited. When the Bayes factor for a model is less than 0.001, the model could be viewed as highly unlikely relative to the other.

Except in a few instances, none of the Bayes factors indicated that one of the alternative scenarios could be considered much less, or more, plausible than the reference case (Table 20). The only scenario with a slightly higher Bayes factor than the reference case was B.1 with

the low prior mean for r . This was consistent with the reference case because the posterior for r in the reference run was updated to support lower values of r than the prior for r .

The two production functions with the B_{msy}/K set at 0.3 and 0.4 had Bayes factors of about 0.2, indicating that the model with B_{msy}/K set at 0.5 provided a somewhat better fit to the data than these alternatives. This also was consistent with the reference run because in spite of the reductions in catch, the abundance indices continue to show a decline in stock size, indicating a highly unproductive stock at low stock sizes and therefore an associated high B_{msy}/K .

The production function with B_{msy}/K set at 0.6 had a Bayes factor of 0.5, also indicating that this model gives a slightly better fit to the data than the lower alternatives. The better fit could be attributed to the Fletcher model's prediction of a sharp drop in surplus production when biomass exceeds B_{msy} when, in contrast, Bocaccio appears to have sustained high exploitation rates for several decades prior to depletion below the B_{msy} level in the 1980s (Figure 17).

The two alternative scenarios which lowered the fixed catch input values for 1986-1995 by factors of 4 and 2 (C.3 and C.4) had the smallest Bayes factors relative to the reference case (0.03 and 0.05, Table 20). This is not surprising because these fixed catches span the period which showed the largest drop in available biomass indices. Consequently, the large drop in biomass levels observed in that period cannot be attributed to a time series of large fixed catches as in the reference case.

Table 20. Bayes factors for alternative mode runs. These reflect the ratio of the probability of the stock assessment data based on a sensitivity run to the probability of the data obtained from the reference case.

| Category Code | Category Description | Code | Run Description | Bayes factor |
|---------------|----------------------|------|--|--------------|
| A | B_{msy}/K | A.1 | $B_{msy}/K = 0.3$ | 0.2 |
| | | A.2 | $B_{msy}/K = 0.4$ | 0.2 |
| | | Ref | $B_{msy}/K = 0.5$ | 1.0 |
| | | A.3 | $B_{msy}/K = 0.6$ | 0.5 |
| B | r prior mean | B.1 | low r (mean = 0.0802, SD = 0.039) | 1.2 |
| | | Ref | reference prior (mean = 0.1067, SD = 0.039) | 1.0 |
| | | B.2 | high r (mean = 0.142, SD = 0.052) | 1.0 |
| C | Catch | C.1 | Sum of trawl and non-halibut HL catch x 0.5, all yrs.' | 0.9 |
| | | Ref. | | 1.0 |
| | | C.2 | Sum of trawl and non-halibut HL catch x 1.5, all yrs.' | 0.3 |
| | | C.3 | Sum of trawl and non-halibut HL catch x 0.25, 86-95 | 0.03 |
| | | C.4 | Sum of trawl and non-halibut HL catch x 0.5, 86-95 | 0.05 |
| F | Catch | F.1 | -2% /y change in Halibut gear q | 0.4 |
| | | F.2 | -1.5% /y change in Halibut gear q | 0.5 |
| | | F.3 | -1.0% /y change in Halibut gear q | 0.6 |
| | | Ref. | 0% /y change in Halibut gear q | 1.0 |
| | | F.4 | 1% /y change in Halibut gear q | 0.8 |
| | | F.5 | 1.5% /y change in Halibut gear q | 0.8 |
| | | F.6 | 2% /y change in Halibut gear q | 0.8 |

13 DECISION TABLES

We have provided forecasting scenarios over 5, 20 (1 generation) and 60 year (3 generations) time horizons for constant catch policies ranging from 0 to 200 t/y for the reference case and 5 sensitivity runs (Table 21). Graphical versions for the reference case are provided in (Figure 25). The forecasts are summarized in the form of decision tables relative to the limit reference point (LRP) and upper target reference point (URP) of $0.4*B_{msy}$ and $0.8*B_{msy}$ respectively (DFO 2006, 2009b), as well as additional relative metrics of stock status.

These projections are based on strong assumptions, including stationarity in model parameters, and that total stock biomass, without reference to the population age or size structure, determines annual surplus production in the following year with no lag. However, these are the same assumptions under which the model reconstruction was made. Therefore, as with most assessments, these long-term projections are provided as guidelines to distinguish between model hypotheses, rather than as true predictions of stock size.

Table 21. Decision tables are provided in the following runs for the reference case and five sensitivity runs.

| Model Run | Decision Table |
|------------------------------|----------------|
| Reference case | Table 22 |
| Case B.1 (low r prior) | Table 23 |
| Case B.2 (high r prior) | Table 24 |
| Case A.1 ($B_{msy}/K=0.3$) | Table 25 |
| Case A.2 ($B_{msy}/K=0.4$) | Table 26 |
| Case A.3 ($B_{msy}/K=0.6$) | Table 27 |

These decision tables are presented to help initiate and focus discussion of harvest strategies for Bocaccio but are not meant to endorse a constant catch policy. Table 29, and Table 30 provide summary decision tables for the probability that stock biomass exceeds 40% of B_{msy} ($0.4*B_{msy}$) within 60 years under each alternative constant TAC policy (t) and under each alternative hypothesized values for r , B_{msy}/K , and historical catch. For example, indicates that, for the reference case, catches of less than 125 t/year are required to have at least a 50% probability of exceeding the LSR point within three generations (60 years). There is some contrast in these projections. For example, the sensitivity run which models $B_{msy}/K=0.3$ predicts that a constant annual harvest of 125 t/year will result in a 0.61 probability of exceeding the LRP in 3 generations while the Reference Case estimates the equivalent probability at 0.49 (Table 29).

Following Edwards et al. (2012), we have also included indicators used by COSEWIC that are based on the decline in the exploitable biomass over 3 generations (i.e., 60 years for Bocaccio) (Table 31). These are COSEWIC indicators A1 and A2 which are used for species that have been assessed as threatened⁷. These indicators are based on the decline in total numbers of mature individuals over the most recent 10 years or 3 generations, whichever is longer, defined as $A1=0.5*N_{t-3Gen}$ (a 50% decline) and $A2=0.7*N_{t-3Gen}$ (a 30% decline), where N_{t-3Gen} is the number of mature individuals three generations previous to year t . However, we used exploitable biomass (B_{t-3Gen}) instead of numbers because of the configuration of the assessment model. Edwards et al. (2012) also present reference points relative to B_0 ($0.2*B_0$ and $0.4*B_0$), which are reference points used by other fishery agencies (e.g. New Zealand Ministry of Fisheries 2007, 2011). However, these have been omitted here because they are identical to the reference points labelled $0.4*B_{msy}$ and $0.8*B_{msy}$, given the Schaefer model assumption that $B_{msy}=0.5*B_0$.

⁷ http://www.cosewic.gc.ca/eng/sct0/assessment_process_e.cfm, updated August 2010

Table 22. Stock status indicators for Bocaccio after 5, 20, and 60 years for the reference case. Policies are constant TAC policies in t . The statistics $P(B > 0.X B_{msy}$ in Hz) refer to the probability that stock size exceeds $0.X B_{msy}$ within the stated horizon (Hz).

| Horizon | Policy | Median(B_{fin}/B_0) | Median(B_{fin}/B_{msy}) | $P(B_{fin} > B_{cur})$ | $P(B > 0.4 * B_{msy}$ in Hz) | $P(B > 0.8 * B_{msy}$ in Hz) |
|----------|--------|-------------------------|-----------------------------|------------------------|---------------------------------|---------------------------------|
| 5 -year | 0 | 0.05 | 0.10 | 0.76 | 0.05 | 0.01 |
| | 50 | 0.05 | 0.09 | 0.67 | 0.05 | 0.01 |
| | 75 | 0.04 | 0.08 | 0.63 | 0.05 | 0.01 |
| | 100 | 0.04 | 0.08 | 0.59 | 0.05 | 0.01 |
| | 125 | 0.04 | 0.07 | 0.53 | 0.04 | 0.01 |
| | 150 | 0.03 | 0.07 | 0.48 | 0.04 | 0.01 |
| | 175 | 0.03 | 0.06 | 0.43 | 0.04 | 0.01 |
| | 200 | 0.03 | 0.06 | 0.37 | 0.04 | 0.01 |
| 20 -year | 0 | 0.16 | 0.33 | 0.88 | 0.43 | 0.21 |
| | 50 | 0.12 | 0.23 | 0.78 | 0.36 | 0.18 |
| | 75 | 0.10 | 0.19 | 0.69 | 0.32 | 0.16 |
| | 100 | 0.07 | 0.15 | 0.61 | 0.28 | 0.14 |
| | 125 | 0.05 | 0.10 | 0.52 | 0.25 | 0.12 |
| | 150 | 0.03 | 0.05 | 0.46 | 0.22 | 0.11 |
| | 175 | 0.01 | 0.02 | 0.41 | 0.19 | 0.10 |
| | 200 | 0.00 | 0.00 | 0.34 | 0.17 | 0.09 |
| 60 -year | 0 | 0.65 | 1.29 | 0.95 | 0.86 | 0.77 |
| | 50 | 0.56 | 1.11 | 0.81 | 0.72 | 0.65 |
| | 75 | 0.44 | 0.88 | 0.69 | 0.65 | 0.58 |
| | 100 | 0.27 | 0.54 | 0.60 | 0.56 | 0.51 |
| | 125 | 0.06 | 0.11 | 0.50 | 0.49 | 0.44 |
| | 150 | 0.00 | 0.000 | 0.44 | 0.42 | 0.38 |
| | 175 | 0.00 | 0.000 | 0.37 | 0.37 | 0.33 |
| | 200 | 0.00 | 0.000 | 0.30 | 0.30 | 0.26 |




Figure 25. Reference case plots of the ratio of a. median stock biomass to B_{msy} for different constant total catch policies and b. 10th, 50th (median), and 90th percentiles.

Table 23. Stock status indicators for Bocaccio after 5, 20, and 60 years for Case B.1 low prior r mean. Policies are constant TAC policies in tons. The statistics $P(B > 0.X * B_{msy}$ in Hz) refer to the probability that stock size exceeds $0.X * B_{msy}$ within the stated horizon (Hz).

| Horizon | Policy | Median(B_{fin}/B_0) | Median(B_{fin}/B_{msy}) | $P(B_{fin} > B_{cur})$ | $P(B > 0.4B_{msy}$ in Hz) | $P(B > 0.8B_{msy}$ in Hz) |
|----------|--------|-------------------------|-----------------------------|------------------------|------------------------------|------------------------------|
| 5 -year | 0 | 0.04 | 0.08 | 0.68 | 0.03 | 0.004 |
| | 50 | 0.04 | 0.08 | 0.58 | 0.03 | 0.004 |
| | 75 | 0.04 | 0.07 | 0.52 | 0.03 | 0.004 |
| | 100 | 0.03 | 0.07 | 0.48 | 0.03 | 0.003 |
| | 125 | 0.03 | 0.06 | 0.44 | 0.03 | 0.003 |
| | 150 | 0.03 | 0.06 | 0.39 | 0.03 | 0.003 |
| | 175 | 0.03 | 0.06 | 0.36 | 0.02 | 0.003 |
| | 200 | 0.03 | 0.05 | 0.32 | 0.02 | 0.003 |
| 20 -year | 0 | 0.10 | 0.19 | 0.81 | 0.29 | 0.13 |
| | 50 | 0.07 | 0.14 | 0.68 | 0.25 | 0.11 |
| | 75 | 0.05 | 0.11 | 0.60 | 0.22 | 0.10 |
| | 100 | 0.04 | 0.07 | 0.51 | 0.20 | 0.09 |
| | 125 | 0.02 | 0.05 | 0.44 | 0.18 | 0.08 |
| | 150 | 0.01 | 0.02 | 0.36 | 0.16 | 0.07 |
| | 175 | 0.00 | 0.00 | 0.31 | 0.14 | 0.06 |
| | 200 | 0.00 | 0.00 | 0.25 | 0.13 | 0.05 |
| 60 -year | 0 | 0.42 | 0.84 | 0.90 | 0.75 | 0.62 |
| | 50 | 0.31 | 0.61 | 0.71 | 0.61 | 0.50 |
| | 75 | 0.18 | 0.35 | 0.60 | 0.54 | 0.43 |
| | 100 | 0.03 | 0.05 | 0.49 | 0.45 | 0.36 |
| | 125 | 0.00 | 0.00 | 0.41 | 0.37 | 0.30 |
| | 150 | 0.00 | 0.000 | 0.33 | 0.31 | 0.26 |
| | 175 | 0.00 | 0.000 | 0.27 | 0.27 | 0.22 |
| | 200 | 0.00 | 0.000 | 0.22 | 0.22 | 0.18 |

Table 24. Stock status indicators for Bocaccio after 5, 20 and 60 years for Case B.2, high prior r mean. Policies are constant TAC policies in tons. The statistics $P(B > 0.X * B_{msy}$ in Hz) refer to the probability that stock size exceeds $0.X * B_{msy}$ within the stated horizon (Hz).

| Horizon | Policy | Median(B_{fin}/B_0) | Median(B_{fin}/B_{msy}) | $P(B_{fin} > B_{cur})$ | $P(B > 0.4B_{msy}$ in Hz) | $P(B > 0.8B_{msy}$ in Hz) |
|----------|--------|-------------------------|-----------------------------|------------------------|------------------------------|------------------------------|
| 5 -year | 0 | 0.05 | 0.10 | 0.80 | 0.06 | 0.011 |
| | 50 | 0.05 | 0.09 | 0.72 | 0.06 | 0.012 |
| | 75 | 0.04 | 0.09 | 0.61 | 0.06 | 0.012 |
| | 100 | 0.04 | 0.08 | 0.56 | 0.06 | 0.011 |
| | 125 | 0.04 | 0.07 | 0.51 | 0.05 | 0.011 |
| | 150 | 0.03 | 0.07 | 0.45 | 0.05 | 0.010 |
| | 175 | 0.03 | 0.06 | 0.40 | 0.05 | 0.010 |
| | 200 | 0.03 | 0.06 | 0.35 | 0.04 | 0.009 |
| 20 -year | 0 | 0.18 | 0.36 | 0.90 | 0.49 | 0.26 |
| | 50 | 0.14 | 0.28 | 0.78 | 0.39 | 0.22 |
| | 75 | 0.10 | 0.19 | 0.71 | 0.36 | 0.20 |
| | 100 | 0.07 | 0.15 | 0.61 | 0.32 | 0.18 |
| | 125 | 0.05 | 0.10 | 0.54 | 0.28 | 0.16 |
| | 150 | 0.02 | 0.05 | 0.46 | 0.25 | 0.15 |
| | 175 | 0.00 | 0.00 | 0.37 | 0.23 | 0.13 |
| | 200 | 0.00 | 0.00 | 0.32 | 0.20 | 0.11 |
| 60 -year | 0 | 0.71 | 1.41 | 0.97 | 0.90 | 0.80 |
| | 50 | 0.61 | 1.22 | 0.81 | 0.76 | 0.68 |
| | 75 | 0.52 | 1.03 | 0.73 | 0.68 | 0.62 |
| | 100 | 0.30 | 0.59 | 0.60 | 0.58 | 0.52 |
| | 125 | 0.10 | 0.20 | 0.52 | 0.50 | 0.46 |
| | 150 | 0.00 | 0.000 | 0.44 | 0.42 | 0.38 |
| | 175 | 0.00 | 0.000 | 0.35 | 0.34 | 0.31 |
| | 200 | 0.00 | 0.000 | 0.28 | 0.29 | 0.26 |

Table 25. Stock status indicators for Bocaccio after 5, 20, and 60 years for Case A.1 B_{msy}/B_0 set at 0.3. Policies are constant TAC policies in tons. The statistics $P(B > 0.X * B_{msy}$ in Hz) refer to the probability that stock size exceeds $0.X * B_{msy}$ within the stated horizon (Hz).

| Horizon | Policy | Median(B_{fin}/B_0) | Median(B_{fin}/B_{msy}) | $P(B_{fin} > B_{cur})$ | $P(B > 0.4B_{msy}$ in Hz) | $P(B > 0.8B_{msy}$ in Hz) |
|----------|--------|-------------------------|-----------------------------|------------------------|------------------------------|------------------------------|
| 5 -year | 0 | 0.04 | 0.14 | 0.76 | 0.13 | 0.03 |
| | 50 | 0.04 | 0.13 | 0.67 | 0.12 | 0.03 |
| | 75 | 0.04 | 0.12 | 0.63 | 0.11 | 0.03 |
| | 100 | 0.03 | 0.12 | 0.59 | 0.11 | 0.03 |
| | 125 | 0.03 | 0.11 | 0.54 | 0.11 | 0.03 |
| | 150 | 0.03 | 0.10 | 0.50 | 0.10 | 0.02 |
| | 175 | 0.03 | 0.10 | 0.46 | 0.10 | 0.02 |
| | 200 | 0.03 | 0.09 | 0.43 | 0.10 | 0.02 |
| 20 -year | 0 | 0.13 | 0.45 | 0.92 | 0.55 | 0.32 |
| | 50 | 0.10 | 0.32 | 0.83 | 0.45 | 0.27 |
| | 75 | 0.08 | 0.27 | 0.74 | 0.42 | 0.26 |
| | 100 | 0.07 | 0.22 | 0.65 | 0.39 | 0.25 |
| | 125 | 0.05 | 0.16 | 0.59 | 0.36 | 0.22 |
| | 150 | 0.03 | 0.11 | 0.50 | 0.34 | 0.21 |
| | 175 | 0.02 | 0.06 | 0.43 | 0.31 | 0.19 |
| | 200 | 0.01 | 0.03 | 0.38 | 0.28 | 0.18 |
| 60 -year | 0 | 0.58 | 1.92 | 0.98 | 0.93 | 0.86 |
| | 50 | 0.46 | 1.54 | 0.92 | 0.86 | 0.78 |
| | 75 | 0.41 | 1.36 | 0.83 | 0.77 | 0.69 |
| | 100 | 0.29 | 0.97 | 0.75 | 0.68 | 0.60 |
| | 125 | 0.21 | 0.69 | 0.68 | 0.61 | 0.53 |
| | 150 | 0.11 | 0.371 | 0.58 | 0.54 | 0.46 |
| | 175 | 0.01 | 0.045 | 0.49 | 0.47 | 0.40 |
| | 200 | 0.01 | 0.033 | 0.41 | 0.40 | 0.35 |

Table 26. Stock status indicators for Bocaccio after 5, 20, and 60 years for Case A.2 B_{msy}/B_0 set at 0.4. Policies are constant TAC policies in tons. The statistics $P(B > 0.X * B_{msy}$ in Hz) refer to the probability that stock size exceeds $0.X * B_{msy}$ within the stated horizon (Hz).

| Horizon | Policy | Median(B_{fin}/B_0) | Median(B_{fin}/B_{msy}) | $P(B_{fin} > B_{cur})$ | $P(B > 0.4B_{msy}$ in Hz) | $P(B > 0.8B_{msy}$ in Hz) |
|----------|--------|-------------------------|-----------------------------|------------------------|------------------------------|------------------------------|
| 5 -year | 0 | 0.05 | 0.11 | 0.74 | 0.10 | 0.02 |
| | 50 | 0.04 | 0.10 | 0.65 | 0.09 | 0.02 |
| | 75 | 0.04 | 0.10 | 0.60 | 0.09 | 0.02 |
| | 100 | 0.04 | 0.09 | 0.54 | 0.09 | 0.02 |
| | 125 | 0.03 | 0.09 | 0.50 | 0.09 | 0.02 |
| | 150 | 0.03 | 0.08 | 0.44 | 0.08 | 0.02 |
| | 175 | 0.03 | 0.07 | 0.40 | 0.07 | 0.02 |
| | 200 | 0.03 | 0.07 | 0.37 | 0.07 | 0.02 |
| 20 -year | 0 | 0.14 | 0.36 | 0.91 | 0.46 | 0.26 |
| | 50 | 0.11 | 0.27 | 0.79 | 0.38 | 0.23 |
| | 75 | 0.09 | 0.22 | 0.73 | 0.35 | 0.22 |
| | 100 | 0.07 | 0.18 | 0.65 | 0.33 | 0.20 |
| | 125 | 0.05 | 0.12 | 0.57 | 0.30 | 0.19 |
| | 150 | 0.03 | 0.08 | 0.52 | 0.28 | 0.16 |
| | 175 | 0.02 | 0.04 | 0.43 | 0.25 | 0.15 |
| | 200 | 0.01 | 0.03 | 0.38 | 0.22 | 0.14 |
| 60 -year | 0 | 0.59 | 1.48 | 0.97 | 0.89 | 0.78 |
| | 50 | 0.48 | 1.19 | 0.90 | 0.79 | 0.68 |
| | 75 | 0.42 | 1.05 | 0.81 | 0.68 | 0.61 |
| | 100 | 0.31 | 0.78 | 0.69 | 0.60 | 0.55 |
| | 125 | 0.19 | 0.47 | 0.60 | 0.54 | 0.49 |
| | 150 | 0.04 | 0.097 | 0.52 | 0.48 | 0.44 |
| | 175 | 0.01 | 0.025 | 0.44 | 0.42 | 0.37 |
| | 200 | 0.01 | 0.025 | 0.37 | 0.36 | 0.32 |

Table 27. Stock status indicators for Bocaccio after 5, 20, and 60 years for Case A.3 B_{msy}/B_0 set at 0.6. Policies are constant TAC policies in tons. The statistics $P(B > 0.X * B_{msy}$ in Hz) refer to the probability that stock size exceeds $0.X * B_{msy}$ within the stated horizon (Hz).

| Horizon | Policy | Median(B_{fin}/B_0) | Median(B_{fin}/B_{msy}) | $P(B_{fin} > B_{cur})$ | $P(B > 0.4B_{msy}$ in Hz) | $P(B > 0.8B_{msy}$ in Hz) |
|----------|--------|-------------------------|-----------------------------|------------------------|------------------------------|------------------------------|
| 5 -year | 0 | 0.06 | 0.10 | 0.78 | 0.10 | 0.02 |
| | 50 | 0.05 | 0.09 | 0.68 | 0.09 | 0.01 |
| | 75 | 0.05 | 0.08 | 0.63 | 0.08 | 0.01 |
| | 100 | 0.05 | 0.08 | 0.56 | 0.08 | 0.01 |
| | 125 | 0.04 | 0.07 | 0.50 | 0.08 | 0.01 |
| | 150 | 0.04 | 0.06 | 0.45 | 0.07 | 0.01 |
| | 175 | 0.04 | 0.06 | 0.39 | 0.07 | 0.01 |
| | 200 | 0.03 | 0.05 | 0.35 | 0.05 | 0.01 |
| 20 -year | 0 | 0.19 | 0.31 | 0.89 | 0.46 | 0.28 |
| | 50 | 0.13 | 0.21 | 0.77 | 0.37 | 0.22 |
| | 75 | 0.09 | 0.15 | 0.69 | 0.34 | 0.20 |
| | 100 | 0.07 | 0.11 | 0.59 | 0.30 | 0.18 |
| | 125 | 0.04 | 0.07 | 0.52 | 0.27 | 0.16 |
| | 150 | 0.02 | 0.03 | 0.42 | 0.23 | 0.14 |
| | 175 | 0.01 | 0.02 | 0.37 | 0.22 | 0.13 |
| | 200 | 0.01 | 0.02 | 0.32 | 0.19 | 0.12 |
| 60 -year | 0 | 0.74 | 1.24 | 0.97 | 0.85 | 0.77 |
| | 50 | 0.62 | 1.03 | 0.84 | 0.73 | 0.65 |
| | 75 | 0.52 | 0.86 | 0.73 | 0.65 | 0.58 |
| | 100 | 0.27 | 0.45 | 0.61 | 0.54 | 0.49 |
| | 125 | 0.06 | 0.09 | 0.52 | 0.47 | 0.43 |
| | 150 | 0.01 | 0.017 | 0.43 | 0.39 | 0.35 |
| | 175 | 0.01 | 0.017 | 0.36 | 0.34 | 0.31 |
| | 200 | 0.01 | 0.017 | 0.31 | 0.30 | 0.27 |

Table 28. Summary decision table for the probability that stock biomass exceeds $0.4*B_{msy}$ within 60 years under each alternative constant TAC policy (t) and under each alternative hypothesized prior mean value for the parameter for the maximum intrinsic rate of increase r .

| | Hypothesized prior mean r | | |
|--------------|-----------------------------|---------------|----------------|
| | Low r (B.1) | Reference r | High r (B.2) |
| Prior mean | 0.0802 | 0.1067 | 0.142 |
| Bayes factor | 1.2 | 1.0 | 1.0 |
| TAC | | | |
| 0 | 0.75 | 0.86 | 0.90 |
| 50 | 0.61 | 0.72 | 0.76 |
| 75 | 0.54 | 0.65 | 0.68 |
| 100 | 0.45 | 0.56 | 0.58 |
| 125 | 0.37 | 0.49 | 0.50 |
| 150 | 0.31 | 0.42 | 0.42 |
| 175 | 0.27 | 0.37 | 0.34 |
| 200 | 0.22 | 0.30 | 0.29 |

Table 29. Summary decision table for the probability that stock biomass exceeds $0.4*B_{msy}$ within 60 years under each alternative constant TAC policy (t) and under each alternative hypothesized value for $t B_{msy}/K$

| B_{msy}/K | Hypothesized B_{msy} to K ratio | | | |
|--------------|-------------------------------------|-----------|------------------|-----------|
| | 0.3 (A.1) | 0.4 (A.2) | Reference 0.5 | 0.6 (A.3) |
| Bayes factor | 0.2 | 0.2 | 1.0 | 0.5 |
| TAC | | | | |
| 0 | 0.93 | 0.89 | 0.86 | 0.85 |
| 50 | 0.86 | 0.79 | 0.72 | 0.73 |
| 75 | 0.77 | 0.68 | 0.65 | 0.65 |
| 100 | 0.68 | 0.60 | 0.56 | 0.54 |
| 125 | 0.61 | 0.54 | 0.49 | 0.47 |
| 150 | 0.54 | 0.48 | 0.42 | 0.39 |
| 175 | 0.47 | 0.42 | 0.37 | 0.34 |
| 200 | 0.40 | 0.36 | 0.30 | 0.30 |

Table 30. Summary decision table for the probability that stock biomass exceeds $0.4*B_{msy}$ within 60 years under each alternative constant TAC policy (t) and under each alternative hypothesized scenario for the level of historic trawl and non-halibut hook and line catch.

| | Hypothesized scenario for historic catch | | |
|----------------|--|------------------|-------------------------------|
| | Low (C.1) 0.5 x ref case | Reference 1.0 | High (C.2) 1.5 x ref. case |
| Catch scenario | 0.5 x ref case | 1.0 | 1.5 x ref. case |
| Bayes factor | 0.9 | 1.0 | 0.3 |
| TAC | | | |
| 0 | 0.82 | 0.86 | 0.89 |
| 50 | 0.62 | 0.72 | 0.82 |
| 75 | 0.52 | 0.65 | 0.68 |
| 100 | 0.40 | 0.56 | 0.61 |
| 125 | 0.30 | 0.49 | 0.54 |
| 150 | 0.23 | 0.42 | 0.49 |
| 175 | 0.18 | 0.37 | 0.42 |
| 200 | 0.14 | 0.30 | 0.37 |

Table 31. Decision table showing the time to reach four reference points (RP): $0.4*B_{msy}$, $0.8*B_{msy}$, $0.5*B_{t-3Gen}$, $0.7*B_{t-3Gen}$ over a range of constant catch quota policies (t) for two levels of confidence for the reference case run (see text for a description of these reference points). Values are the first year that the RP is reached with the given confidence level (and the population is increasing). Declining outcomes were found for more policies under the 80% confidence level for the B_0 and B_{msy} reference points since at 80% the intervals get wider on smaller quota policies than for the medians.

| Quota Policy | | | | |
|----------------|---------------|---------------|------------------|------------------|
| 50% Confidence | $0.4*B_{msy}$ | $0.8*B_{msy}$ | $0.5*B_{t-3Gen}$ | $0.7*B_{t-3Gen}$ |
| 0 | 23 | 41 | 21 | 24 |
| 50 | 30 | 49 | 24 | 31 |
| 75 | 37 | 58 | 30 | 34 |
| 100 | 46 | >60 | 34 | 37 |
| 125 | >60 | >60 | 41 | 43 |
| 150 | declining | declining | declining | declining |
| 175 | declining | declining | declining | declining |
| 200 | declining | declining | declining | declining |
| 80% Confidence | | | | |
| 0 | 59 | >60 | 36 | 38 |
| 50 | >60 | >60 | 43 | 52 |
| 75 | declining | declining | >60 | >60 |
| 100 | declining | declining | >60 | >60 |
| 125 | declining | declining | >60 | >60 |
| 150 | declining | declining | declining | declining |
| 175 | declining | declining | declining | declining |
| 200 | declining | declining | declining | declining |

14 STATUS OF BOCACCIO IN U.S. WATERS

Only the California and southern Oregon portion of the U.S. population of Bocaccio has been assessed in recent years. The most recent assessment was provided in November 2011, but is still only available in draft form (Field 2011). Field (2011) reports that the results are slightly more pessimistic relative to the 2009 model, with depletion of spawning biomass in the year 2011 estimated at 26% of B_0 relative to the 30% projected from the 2009 model. Continued decline in the trawl survey and hook and line survey indices were mainly responsible for this change. A young-of-the-year index suggests a flattening of what was previously an increasing trend.

Field (2011) notes further that spawning output [estimated biomass in egg production] exhibits a very moderate decline until about 1950, with a steep decline from the early 1950s followed by a sharp increase in the early 1960s. Spawning output is estimated to have exceeded the mean unfished biomass level through the early 1970s, when high fishing mortality rates again resulted in a rapid decline. Harvests declined towards the end of the 1990s, in response to management restrictions. Since the early 2000s, spawning output has been increasing steadily, largely as a result of reduced fishing mortality and a strong 1999 year class, although the rate of increase has slowed in the later half of the 2000s. Indications of strong 2009 and 2010 year classes should lead to additional increases in abundance.




Figure 26. Estimated spawning output time series 1892-2011 for the base case fir California and southern Oregon Bocaccio population, with approximate 95% confidence intervals (figure from Field 2011).

15 SUMMARY

This document provides a stock assessment for Bocaccio in BC waters. Results of the work are intended to serve as advice over the short term to managers and stakeholders on stock status, and likely impacts of different fixed harvest options.

The reference case analysis indicates that is likely that the Bocaccio population in BC has been declining for many decades and is currently well below the LRP of $0.4*B_{msy}$. Furthermore, while there is considerable uncertainty in estimating current trends, there is no sign that the population has started to increase, and appears to have continued to decline over the most recent decade. Current harvests are approximately equal to estimates of replacement yield. The impacts on estimates of stock status of alternative model assumptions to those made in the reference case were explored with additional sensitivity runs. These runs were, in general terms, consistent with the reference case results.

Long term biomass projections were made for the reference case and a selection of the sensitivity runs over 5, 20, and 60 year scenarios under varying fixed harvest assumptions. These projections are shown relative to the DFO draft policy target reference points of $0.4*B_{msy}$ and $0.8*B_{msy}$ and other reference points.

While the Bayesian approach used in this assessment provides a formal mechanism to include uncertainty in model output (including predictions), managers, and stakeholders are advised that not all sources of uncertainty have been addressed and that it is likely that the true uncertainty is even greater than that presented here.

16 RESPONSES TO 2009 RECOMMENDATIONS FOR FUTURE WORK

The following section summarizes the authors' responses to recommendations (in italics) made during review of the earlier work (Stanley et al. 2009a).

1. *Consider using the number of troll licenses as a surrogate for relative troll effort in the reconstruction of bycatch in the early salmon troll fishery.*

The authors recognize that additional work could go into the catch reconstructions for each sector. However, the model is not particularly sensitive to modest changes in historical catches. Changes in pre-1950 troll catches of Bocaccio would have little impact. Finally, there are a large number of alternative means for reconstructing catches with little objective basis for choosing amongst them.

2. *Explore the potential to work with US biologists for a coastwide assessment of Bocaccio, especially as the time series of abundance indices and ageing data expands.*

This was not yet examined. While US assessments have so far concentrated on California data, US staff have expressed, as well, a desire to do more collaborative work, especially with ageing. Canadian and US staff have been collaborating on Bocaccio genetics work. Canadian samples were included in the genetics work noted above.

3. *Develop software and an empirical basis to carry out management strategy evaluation (MSE) of alternative feedback control fisheries management regimes for Bocaccio alone or combinations of rockfish species.*

Some preliminary work on MSE work has been conducted on Bocaccio⁸. This work focussed on whether the current surveys can provide adequate monitoring of Bocaccio abundance. The unpublished work indicated that that in spite of the imprecision of each survey, when considered collectively in a modelling context, they could provide adequate monitoring. No further MSE work has been conducted or is planned.

4. *Examine the feasibility of a trolling or gillnet experiment to estimate the ratio of the densities of Bocaccio or other species in trawlable and untrawlable areas.*

As noted above, results from Matthews et al. (1989) gillnet survey were used in this assessment.

5. *Update the model to address the reviewer's suggestion that the model account for the fact that a significant portion of the area within each trawlable block may, in fact, be untrawlable.*

The assessment did not incorporate this 2009 reviewer's comment. However, we note that a significant portion of the area within each untrawlable block may, in fact, be trawlable which would act to compensate. We have no information on these two proportions.

6. *Evaluate the possibility of obtaining additional prior information of the survey net catchability coefficient by studying the relationship between stock size estimates and groundfish survey area swept estimates in the U.S. Bocaccio assessments.*

Sufficient time was not available for the authors to consider incorporating U.S. Bocaccio survey catchability in this U.S. assessment. The use of different vessels, nets, and different bottom type, would imply that the values would not be comparable; however, the comparison could be informative.

7. *Evaluate the feasibility of a stock structure study of Bocaccio in BC and US waters using samples of chemical microconstituents in Bocaccio body parts. The presence of much older fish in recent samples from BC and Washington State in comparison with California*

⁸ McAllister, M.K., Stanley, R., and Kronlund, R. 2009. Can trawl surveys tell us whether a recovery plan is working? Poster presented at ICES/PICES/UNCOVER Symposium on Rebuilding Depleted Fish Stocks –Biology, Ecology, Social Science and Management Strategies. Warnemünde/Rostock, Germany.

samples, in spite of significant fishing mortality for many decades, implies the possibility of gradual migration to BC waters as US fish become older. Microconstituent analysis might reveal the source of larvae and juveniles that recruit to BC fisheries.

Resources and time were not sufficient available to conduct microconstituent analysis of Bocaccio samples.

8. *Evaluate the feasibility of acoustic studies of Bocaccio or other rockfish behaviour in response to trawl gear.*

No rockfish acoustic studies were conducted. Rockfish acoustic work is problematic for a variety of reasons including the difficulty in identify the rockfish to species and the difficulty in using ship-based acoustics on near bottom targets, within the acoustic “dead-zone”.

9. *Examine post-model pre-data distribution of model outputs.*

This procedure was adopted and summarized above.

17 RECOMMENDATIONS FOR FUTURE WORK

Subject to the availability of research resources and the many other competing priorities related to the more than 100 other exploited populations of groundfish on the Pacific coast of Canada, we suggest that consideration be given to the following research directions:

1. Continue to work with U.S. biologists on Bocaccio research issues and, if possible, a coastwide assessment of Bocaccio.
2. Publish the nearly completed work on an MSE-based study of the adequacy of the current survey array in tracking Bocaccio abundance.
3. For the next assessment, consider incorporating end-of-summer YOY length-at-age data (Russ Markell, pers. comm., University of British Columbia) in estimation for growth parameters.
4. Conduct a review of Bocaccio surveys trends in 5 years to check for evidence of further declines in abundance and, if appropriate include results of the DFO longline surveys in this review and subsequent assessments.
5. Conduct a full assessment of Bocaccio in approximately 2022. The timing will coincide with an anticipated COSEWIC assessment.
6. We recommend continued sampling and ageing of Bocaccio. However, we note the limited amount of ageing resources and the large number of groundfish species/populations to be assessed. It might be advantageous to direct ageing resources to species for which representative time series can be developed or that currently lack sufficient material to estimate life history parameters.

ACKNOWLEDGEMENTS

We thank Dr. Alexandra Valentin for bringing to our attention recent genetics work as well as the US identification of a separate population for Puget Sound/Georgia Basin. We also thank Norm Olsen and Kate Rutherford for assisting in extracting catch, survey, and biological data from DFO data bases and assisting with the tables and figures.

LITERATURE CITED

- Boutillier, J.A. 1998. Queen Charlotte Sound shrimp survey and resulting management actions, July 1998. Can. Man. Rep. Fish. Aquat. Sci. 2462.
- Bell, F.H., H.A. Dunlop, and Freeman, N.L. 1952. Pacific coast halibut landings, 1888-1950 and catch according to area of origin. Rep. Int. Fish. Comm. 17.
- Buonaccorsi, V.P., Kimbrell, C.A., Lynn, E.A., and Hyde, J.R. 2012. Comparative population genetic analysis of Bocaccio rockfish *Sebastes paucispinis* using anonymous and gene-associated simple sequence repeat loci. J. Hered. 103(3):391-399.
- Buonaccorsi, V.P., Starkey, E., and Graves, J.E. 2001. Mitochondrial and nuclear DNA analysis of population subdivision among young-of-the-year Spanish mackerel (*Scomberomorus maculatus*) from the western Atlantic and Gulf of Mexico. Mar. Biol. 138:37-45
- COSEWIC. 2002. COSEWIC Status Report – Bocaccio.
http://www.sararegistry.gc.ca/document/default_e.cfm?documentID=174.
- DFO. 2006. A harvest strategy compliant with a precautionary approach. DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2006/023.
- DFO. 2009a. Recovery Potential Assessment for Bocaccio (*Sebastes paucispinis*). DFO Can. Sci. Advis. Sec. Sci. Advis. Rep. 2009/040.
- DFO. 2009b. A fishery decision-making framework incorporating the Precautionary Approach. Last modified 23rd March 2009. <http://www.dfo-mpo.gc.ca/fm-gp/peches-fisheries/fish-ren-peche/sff-cpd/precaution-eng.htm>
- Drake, J.S., Berntson, E.A., Cope, J.M., Gustafson, R.G., Holmes, E.E., Levin, P.S., Tolimieri, N., Waples, R.S., Sogard, S.M., and Williams, G.D. 2010. Status review of five rockfish species in Puget Sound, Washington: Bocaccio (*Sebastes paucispinis*), canary rockfish (*S. pinniger*), yelloweye rockfish (*S. ruberrimus*), greenstriped rockfish (*S. elongatus*), and redstripe rockfish (*S. proriger*). U.S. Dept. Commer., NOAA Tech. Memo. NMFS-NWFSC-108, 234 p.
- Edwards, A.M., Haigh, R., and Starr, P.J. 2012. Stock assessment and recovery potential assessment for Yellowmouth Rockfish (*Sebastes reedi*) along the Pacific coast of Canada. DFO Can. Sci. Advis. Sec. Res. Doc. 2012/095. iv +188p. http://www.dfo-mpo.gc.ca/csas-sccs/Publications/ResDocs-DocRech/2012/2012_095-eng.html
- Efron, B. 1982. The jackknife, the bootstrap, and other resampling plans. Society for Industrial and Applied Mathematics. Philadelphia.
- Field, J. 2011. Status of Bocaccio, *Sebastes paucispinis*, in the Conception, Monterey and Eureka INPFC areas as evaluated for 2011. http://www.pcouncil.org/wp-content/uploads/E1a_ATT1_STATUS_BOCACCIO_NOV2011BB.pdf.
- Field, J.C., Dick, E.J., Pearson, D., and MacCall, A.D. 2009. Status of Bocaccio, *Sebastes paucispinis*, in the Conception, Monterey and Eureka INPFC areas for 2009. http://www.pcouncil.org/wp-content/uploads/Bocaccio_Final_Jan15_2010.pdf
- Forrest, R.E., McAllister, M.K., Dorn, M.W., Martell, S.J.D., and Stanley, R.D. 2010. Hierarchical Bayesian estimation of recruitment parameters and reference points for Pacific rockfishes (*Sebastes* spp.) under alternative assumptions about the stock–recruit function. Can. J. Fish. Aquat. Sci. 67: 1611–1634.
- Hilborn, R. and Walters, C.J. 1992. Quantitative fisheries stock assessment: choice, dynamics. Routledge, Chapman and Hall. New York.
- Kass, R.E. and Raftery, A.E. 1995. Bayes factors. J. Am. Stat. Assoc. 90: 773–795.

-
- King, J.R., McAllister, M., Holt, K.R. and Starr, P.J. 2012. Lingcod (*Ophiodon elongatus*) stock assessment and yield advice for outside stocks in British Columbia. DFO Can. Sci. Advis. Sec. Res. Doc. 2011/124. viii + 177p.
- Love, M.S., Yoklavich, M., and Thorsteinsson, L. 2002. University of California Press. London.
- Matthews, K.R., Candy, J.R., Richards, L.J., and Hand, C.M. 1989. Experimental gillnet fishing on trawlable and untrawlable areas off Northwestern Vancouver Island, from the MV Caledonian August 15-28, 1989. Can. Man. Rep. Fish. Aquat. Sci. No. 2046.
- McAllister, M.K. and Babcock, E.A. 2002. Bayesian surplus production model with the Sampling Importance Resampling algorithm (BSP): a user's guide. National Marine Fisheries Service 2002 Shark Evaluation Workshop, Panama City, Florida. SB-02-26. Also see: BSP (3.0) at: <http://www.iccat.int/en/AssessCatalog.htm>
- McAllister, M.K. and Kirchner, C.H. 2002. Accounting for structural uncertainty to facilitate precautionary fishery management: illustration with Namibian orange roughy. Bull. Mar. Sci. 70(2): 499-540.
- McAllister, M.K. and Kirkwood, G.P. 1998. Using Bayesian decision analysis to help achieve a precautionary approach to managing newly developing fisheries. Can. J. Fish. Aquat. Sci. 55: 2642-2661.
- McAllister, M.K., Babcock, E.A., and Pikitch, E.K. 2001a. Using Bayesian methods to provide an empirical basis to weight conflicting stock assessment results on stock rebuilding when there is uncertainty over processes affecting future recruitment. ICCAT Col. Vol. Sci. Papers. SCRS/00/103.
- McAllister, M.K., Pikitch, E.K., and Babcock, E.A. 2001b. Using demographic methods to construct Bayesian priors for the intrinsic rate of increase in the Schaefer model and implications for stock rebuilding. Can. J. Fish. Aquat. Sci. 58(9): 1871-1890.
- McAllister, M.K., Stanley, R.D., and Starr, P. 2010. Modeling trawl survey catchability for BC bocaccio (*Sebastes paucispinis*): a combined expert driven – empirical approach. Fish. Bull. 108:282-304
- McAllister, M.K., Babcock, E.A., Pikitch, E.K., and Prager, M.H. 1999. Application of a non-equilibrium generalized production model to South and North Atlantic Swordfish: combining Bayesian and demographic methods for parameter estimation. ICCAT Col. Vol. Sci. Papers. SCRS/99/85.
- Meyer, R. and Millar, R.B. 1999. BUGS in Bayesian stock assessments. Can. J. Fish. Aquat. Sci. 56: 1078-1086.
- New Zealand Ministry of Fisheries. 2007. Operational guidelines for New Zealand's harvest strategy standard. 67 p. <http://www.fish.govt.nz/NR/rdonlyres/8419AC9A-8DDA-48A6-833C-AF6E70052244/0/OperationalGuidelinesHSS.pdf>
- New Zealand Ministry of Fisheries. 2011. Operational guidelines for New Zealand's harvest strategy standard. June 2011, 78 p. http://fs.fish.govt.nz/Doc/22847/Operational_Guidelines_for_HSS_rev_1_Jun_2011.pdf.ashx
- Olsen, N., Workman, G.D. and Stanley, R.D. 2007. Queen Charlotte Sound bottom trawl survey, July 5th to August 9th, 2005. Can. Man. Rep. Fish. Aquat. Sci. 2784.
- Prager, M.H. 1994. A suite of extensions to a nonequilibrium surplus production model. Fish. Bull. 92: 374-389.
-

-
- Punt, A.E. and Butterworth, D.S. 2002. An examination of certain of the assumptions made in the Bayesian approach used to assess the eastern north Pacific stock of gray whales (*Eschrichtius robustus*). *J. Cetacean Res. Manage.* 4(1):99-110.
- Quinn, T.R. and Deriso, R.B. 1999. *Quantitative Fish Dynamics*. Oxford University Press.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. *Bull. Fish. Brd. Can.* 191.
- Stanley, R.D. and Kronlund, A.R. 2005. Life history characteristics for silvergray rockfish (*Sebastes brevispinis*) in British Columbia waters and the implications for stock assessment and management. *Fish. Bull.* 103: 670-684.
- Stanley, R.D., McAllister, M. Starr, P., and Olsen, N. 2009a. Stock assessment for bocaccio (*Sebastes paucispinis*) in British Columbia waters. *DFO Can. Sci. Advis. Sec. Res. Doc.* 2009/055.
- Stanley, R.D., Olsen, N., and Fedoruk, A. 2009b. The accuracy of yelloweye rockfish catch estimates from the British Columbia Groundfish Integration Project. *Marine and Coastal Fisheries: Dynamics, Management, and Ecosystem Science* 1:354–362.
- Starr, P.J., Sinclair, A.S., and Boutillier, J. 2002. West Coast Vancouver Island Pacific cod assessment. *DFO. Can. Sci. Advis. Sec. Res. Doc.* 2002/113.
- Weinberg, K.L., Wilkins, M.E., Shaw, F.R., and Zimmerman, M. 2001. The 2001 Pacific west coast bottom trawl survey of groundfish resources: estimates of distribution, abundance, and length and age composition. *NOAA Technical Memorandum NMFS-AFSC-128*.
- Wishard, L.N., Utter, F.M., and Gunderson, D.R. 1980. Stock separation of five rockfish species using naturally occurring biochemical genetic markers. *Mar. Fish. Rev.* 42:64-73.
- Workman, G.D., Olsen, N., and Rutherford, K.L. 2007. West Coast Charlotte Islands groundfish bottom trawl survey, August 28th to September 25th, 2006. *Can. Man. Rep. Fish. Aquat. Sci.* 2840
- Workman, G.D., Rutherford, K.L., and Olsen, N. 2008a. Hecate Strait groundfish bottom trawl survey, May 25 to June 29, 2005. *Can. Man. Rep. Fish. Aquat. Sci.* 2805.
- Workman, G.D., Olsen, N., Fargo, J.J., and Stanley, R.D. 2008b. West Coast Vancouver Island groundfish bottom trawl survey, R/V W.E.Ricker, May 25 to June 12, 2004. *Can. Man. Rep. Fish. Aquat. Sci.* 2826.
- Yamanaka, K.L., McAllister, M.K., Etienne, M-P., and Flemming, R. 2012. Stock Assessment and Recovery Potential Assessment for Quillback Rockfish (*Sebastes maliger*) on the Pacific Coast of Canada. *DFO Can. Sci. Advis. Sec. Res. Doc.* 2011/135. vii + 151p.

APPENDIX A CATCH

Appendix Table 1. Inputted catch values in the Reference case (1935-1975). Catches for Trawl and ZN HL are fixed; catches in the other fisheries are estimated. Note that catch values are rounded to nearest ton so percentages do not exactly match.

| Year | Fixed | Estimated (Medians) | | | Total | Year | Proportion of Total Catch (%) | | | |
|------|-----------------|---------------------|--------------|-------|-------|-----------------|-------------------------------|--------------|-------|----|
| | Trawl and ZN HL | Halibut | Salmon Troll | Recr. | | Trawl and ZN HL | Halibut | Salmon Troll | Recr. | |
| 1935 | 1 | 1242 | 393 | 3 | 1639 | 1935 | 0% | 76% | 24% | 0% |
| 1936 | 1 | 1360 | 381 | 3 | 1745 | 1936 | 0% | 78% | 22% | 0% |
| 1937 | 1 | 1199 | 365 | 3 | 1568 | 1937 | 0% | 76% | 23% | 0% |
| 1938 | 2 | 1043 | 347 | 3 | 1396 | 1938 | 0% | 75% | 25% | 0% |
| 1939 | 2 | 1237 | 312 | 3 | 1554 | 1939 | 0% | 80% | 20% | 0% |
| 1940 | 11 | 1212 | 290 | 3 | 1516 | 1940 | 1% | 80% | 19% | 0% |
| 1941 | 8 | 1121 | 401 | 3 | 1532 | 1941 | 1% | 73% | 26% | 0% |
| 1942 | 36 | 981 | 377 | 3 | 1397 | 1942 | 3% | 70% | 27% | 0% |
| 1943 | 100 | 948 | 489 | 2 | 1540 | 1943 | 6% | 62% | 32% | 0% |
| 1944 | 45 | 723 | 145 | 2 | 915 | 1944 | 5% | 79% | 16% | 0% |
| 1945 | 418 | 701 | 317 | 2 | 1438 | 1945 | 29% | 49% | 22% | 0% |
| 1946 | 213 | 804 | 246 | 3 | 1265 | 1946 | 17% | 64% | 19% | 0% |
| 1947 | 116 | 700 | 396 | 5 | 1218 | 1947 | 10% | 58% | 33% | 0% |
| 1948 | 183 | 690 | 277 | 8 | 1158 | 1948 | 16% | 60% | 24% | 1% |
| 1949 | 221 | 666 | 385 | 10 | 1282 | 1949 | 17% | 52% | 30% | 1% |
| 1950 | 209 | 677 | 411 | 12 | 1309 | 1950 | 16% | 52% | 31% | 1% |
| 1951 | 200 | 795 | 430 | 14 | 1439 | 1951 | 14% | 55% | 30% | 1% |
| 1952 | 187 | 754 | 339 | 16 | 1296 | 1952 | 14% | 58% | 26% | 1% |
| 1953 | 78 | 550 | 336 | 18 | 982 | 1953 | 8% | 56% | 34% | 2% |
| 1954 | 81 | 566 | 291 | 20 | 959 | 1954 | 8% | 59% | 30% | 2% |
| 1955 | 104 | 472 | 356 | 22 | 954 | 1955 | 11% | 49% | 37% | 2% |
| 1956 | 98 | 469 | 334 | 23 | 923 | 1956 | 11% | 51% | 36% | 2% |
| 1957 | 74 | 525 | 372 | 26 | 997 | 1957 | 7% | 53% | 37% | 3% |
| 1958 | 70 | 494 | 364 | 28 | 955 | 1958 | 7% | 52% | 38% | 3% |
| 1959 | 91 | 538 | 354 | 29 | 1013 | 1959 | 9% | 53% | 35% | 3% |
| 1960 | 66 | 484 | 358 | 30 | 938 | 1960 | 7% | 52% | 38% | 3% |
| 1961 | 92 | 463 | 393 | 33 | 980 | 1961 | 9% | 47% | 40% | 3% |
| 1962 | 164 | 491 | 344 | 31 | 1030 | 1962 | 16% | 48% | 33% | 3% |
| 1963 | 144 | 541 | 311 | 31 | 1028 | 1963 | 14% | 53% | 30% | 3% |
| 1964 | 110 | 427 | 330 | 31 | 898 | 1964 | 12% | 48% | 37% | 3% |
| 1965 | 290 | 389 | 347 | 32 | 1058 | 1965 | 27% | 37% | 33% | 3% |
| 1966 | 1073 | 343 | 312 | 29 | 1757 | 1966 | 61% | 20% | 18% | 2% |
| 1967 | 785 | 315 | 344 | 28 | 1472 | 1967 | 53% | 21% | 23% | 2% |
| 1968 | 533 | 284 | 359 | 27 | 1204 | 1968 | 44% | 24% | 30% | 2% |
| 1969 | 1064 | 359 | 315 | 26 | 1765 | 1969 | 60% | 20% | 18% | 1% |
| 1970 | 457 | 304 | 294 | 26 | 1081 | 1970 | 42% | 28% | 27% | 2% |
| 1971 | 324 | 255 | 311 | 27 | 917 | 1971 | 35% | 28% | 34% | 3% |
| 1972 | 452 | 283 | 274 | 28 | 1038 | 1972 | 44% | 27% | 26% | 3% |
| 1973 | 1112 | 196 | 234 | 28 | 1569 | 1973 | 71% | 12% | 15% | 2% |
| 1974 | 1274 | 131 | 233 | 27 | 1665 | 1974 | 77% | 8% | 14% | 2% |

Appendix Table 2. Inputted catches in the Reference case (1976-2012). Catches for Trawl and ZN HL are fixed; catches in the other fisheries are estimated. Note that catch values are rounded to nearest ton so percentages do not exactly match

| Year | Fixed Trawl and ZN HL | Estimated (Medians) | | | Total | Year | Proportion of Total Catch (%) | | | |
|------|--------------------------------|---------------------|-----------------|-------|-------|------|-------------------------------|---------|-----------------|-------|
| | | Halibut | Salmon Troll | Recr. | | | Trawl and ZN HL | Halibut | Salmon Troll | Recr. |
| 1975 | 790 | 184 | 224 | 26 | 1224 | 1975 | 65% | 15% | 18% | 2% |
| 1976 | 677 | 233 | 220 | 27 | 1157 | 1976 | 59% | 20% | 19% | 2% |
| 1977 | 399 | 151 | 228 | 28 | 807 | 1977 | 49% | 19% | 28% | 4% |
| 1978 | 255 | 127 | 273 | 30 | 684 | 1978 | 37% | 19% | 40% | 4% |
| 1979 | 486 | 173 | 274 | 30 | 962 | 1979 | 51% | 18% | 28% | 3% |
| 1980 | 183 | 149 | 364 | 31 | 726 | 1980 | 25% | 21% | 50% | 4% |
| 1981 | 95 | 144 | 310 | 31 | 580 | 1981 | 16% | 25% | 53% | 5% |
| 1982 | 105 | 138 | 304 | 32 | 580 | 1982 | 18% | 24% | 53% | 6% |
| 1983 | 154 | 131 | 282 | 32 | 599 | 1983 | 26% | 22% | 47% | 5% |
| 1984 | 176 | 98 | 260 | 33 | 566 | 1984 | 31% | 17% | 46% | 6% |
| 1985 | 418 | 115 | 252 | 28 | 814 | 1985 | 51% | 14% | 31% | 3% |
| 1986 | 720 | 134 | 196 | 15 | 1065 | 1986 | 68% | 13% | 18% | 1% |
| 1987 | 732 | 120 | 155 | 25 | 1032 | 1987 | 71% | 12% | 15% | 2% |
| 1988 | 1348 | 102 | 141 | 16 | 1607 | 1988 | 84% | 6% | 9% | 1% |
| 1989 | 808 | 79 | 123 | 22 | 1033 | 1989 | 78% | 8% | 12% | 2% |
| 1990 | 1063 | 43 | 136 | 21 | 1263 | 1990 | 84% | 3% | 11% | 2% |
| 1991 | 1093 | 37 | 116 | 22 | 1268 | 1991 | 86% | 3% | 9% | 2% |
| 1992 | 976 | 28 | 106 | 24 | 1134 | 1992 | 86% | 3% | 9% | 2% |
| 1993 | 1160 | 25 | 66 | 14 | 1266 | 1993 | 92% | 2% | 5% | 1% |
| 1994 | 635 | 20 | 44 | 15 | 714 | 1994 | 89% | 3% | 6% | 2% |
| 1995 | 545 | 16 | 31 | 9 | 601 | 1995 | 91% | 3% | 5% | 2% |
| 1996 | 343 | 15 | 17 | 4 | 378 | 1996 | 91% | 4% | 4% | 1% |
| 1997 | 267 | 18 | 12 | 9 | 306 | 1997 | 87% | 6% | 4% | 3% |
| 1998 | 236 | 19 | 7 | 10 | 273 | 1998 | 86% | 7% | 3% | 4% |
| 1999 | 251 | 20 | 4 | 11 | 286 | 1999 | 88% | 7% | 1% | 4% |
| 2000 | 303 | 16 | 3 | 8 | 330 | 2000 | 92% | 5% | 1% | 2% |
| 2001 | 288 | 15 | 3 | 8 | 313 | 2001 | 92% | 5% | 1% | 3% |
| 2002 | 295 | 17 | 7 | 9 | 328 | 2002 | 90% | 5% | 2% | 3% |
| 2003 | 237 | 16 | 8 | 10 | 270 | 2003 | 88% | 6% | 3% | 4% |
| 2004 | 170 | 17 | 9 | 9 | 205 | 2004 | 83% | 8% | 4% | 4% |
| 2005 | 162 | 18 | 12 | 9 | 201 | 2005 | 81% | 9% | 6% | 4% |
| 2006 | 131 | 16 | 11 | 9 | 167 | 2006 | 79% | 10% | 7% | 5% |
| 2007 | 139 | 13 | 8 | 7 | 166 | 2007 | 84% | 8% | 5% | 4% |
| 2008 | 118 | 11 | 5 | 7 | 140 | 2008 | 84% | 8% | 4% | 5% |
| 2009 | 114 | 8 | 6 | 6 | 134 | 2009 | 85% | 6% | 4% | 5% |
| 2010 | 99 | 7 | 6 | 6 | 118 | 2010 | 84% | 6% | 5% | 5% |
| 2011 | 119 | 6 | 6 | 6 | 137 | 2011 | 87% | 4% | 4% | 4% |
| 2012 | 119 | 6 | 6 | 6 | 137 | 2012 | 87% | 4% | 4% | 5% |

Appendix Table 3. Time series of fishery effort in the halibut, salmon troll, and recreational fisheries

| Year | Halibut fishery effort | Salmon troll fishery effort | Recreational fishery effort | Catch of Bocaccio in halibut fishery | Year | Halibut fishery effort | Salmon troll fishery effort | Recreational fishery effort | Catch of Bocaccio in halibut fishery |
|------|------------------------|-----------------------------|-----------------------------|--------------------------------------|------|------------------------|-----------------------------|-----------------------------|--------------------------------------|
| | (100,000 skates) | (10,000 boat days) | (100,000 angler days) | (mt) | | (100,000 skates) | (10,000 boat days) | (100,000 angler days) | (mt) |
| 1935 | 2.35 | 3.57 | 0.17 | NA | 1975 | 1.05 | 7.14 | 4.44 | NA |
| 1936 | 2.70 | 3.57 | 0.17 | NA | 1976 | 1.37 | 7.21 | 4.61 | NA |
| 1937 | 2.51 | 3.57 | 0.17 | NA | 1977 | 0.89 | 7.35 | 4.79 | NA |
| 1938 | 2.26 | 3.57 | 0.17 | NA | 1978 | 0.73 | 8.63 | 4.97 | NA |
| 1939 | 2.83 | 3.32 | 0.17 | NA | 1979 | 1.01 | 8.80 | 5.15 | NA |
| 1940 | 2.79 | 3.25 | 0.17 | NA | 1980 | 0.87 | 11.90 | 5.32 | NA |
| 1941 | 2.65 | 4.67 | 0.17 | NA | 1981 | 0.84 | 9.97 | 5.49 | NA |
| 1942 | 2.39 | 4.55 | 0.17 | NA | 1982 | 0.81 | 9.96 | 5.67 | NA |
| 1943 | 2.32 | 6.11 | 0.17 | NA | 1983 | 0.80 | 9.31 | 5.84 | NA |
| 1944 | 1.77 | 1.88 | 0.17 | NA | 1984 | 0.62 | 9.13 | 6.01 | NA |
| 1945 | 1.75 | 4.27 | 0.17 | NA | 1985 | 0.75 | 9.21 | 5.62 | NA |
| 1946 | 2.12 | 3.35 | 0.20 | NA | 1986 | 0.97 | 7.80 | 3.20 | NA |
| 1947 | 1.96 | 5.60 | 0.41 | NA | 1987 | 0.96 | 6.82 | 5.82 | NA |
| 1948 | 1.96 | 3.99 | 0.60 | NA | 1988 | 0.94 | 6.99 | 4.36 | NA |
| 1949 | 1.92 | 5.70 | 0.80 | NA | 1989 | 0.79 | 6.51 | 6.69 | NA |
| 1950 | 1.99 | 6.15 | 1.00 | NA | 1990 | 0.49 | 8.17 | 7.37 | NA |
| 1951 | 2.48 | 6.74 | 1.21 | NA | 1991 | 0.49 | 8.02 | 8.53 | NA |
| 1952 | 2.34 | 5.44 | 1.40 | NA | 1992 | 0.45 | 8.55 | 11.15 | NA |
| 1953 | 1.75 | 5.48 | 1.60 | NA | 1993 | 0.52 | 6.96 | 8.24 | NA |
| 1954 | 1.78 | 4.74 | 1.80 | NA | 1994 | 0.46 | 5.24 | 9.99 | NA |
| 1955 | 1.51 | 5.79 | 2.01 | NA | 1995 | 0.44 | 4.20 | 7.08 | NA |
| 1956 | 1.54 | 5.59 | 2.20 | NA | 1996 | 0.42 | 2.38 | 2.81 | NA |
| 1957 | 1.75 | 6.35 | 2.40 | NA | 1997 | 0.52 | 1.80 | 7.27 | NA |
| 1958 | 1.71 | 6.31 | 2.61 | NA | 1998 | 0.57 | 1.05 | 8.43 | NA |
| 1959 | 1.81 | 6.20 | 2.81 | NA | 1999 | 0.60 | 0.58 | 8.82 | NA |
| 1960 | 1.66 | 6.33 | 3.00 | NA | 2000 | 0.47 | 0.47 | 6.11 | NA |
| 1961 | 1.60 | 7.21 | 3.38 | NA | 2001 | 0.46 | 0.38 | 6.66 | NA |
| 1962 | 1.70 | 6.40 | 3.38 | NA | 2002 | 0.54 | 1.20 | 7.93 | NA |
| 1963 | 1.84 | 5.84 | 3.38 | NA | 2003 | 0.54 | 1.29 | 9.07 | NA |
| 1964 | 1.50 | 6.35 | 3.38 | NA | 2004 | 0.60 | 1.50 | 8.70 | NA |
| 1965 | 1.40 | 6.89 | 3.38 | NA | 2005 | 0.63 | 2.09 | 8.54 | NA |
| 1966 | 1.31 | 6.60 | 3.38 | NA | 2006 | 0.60 | 1.97 | 8.72 | 8.09 |
| 1967 | 1.24 | 7.57 | 3.38 | NA | 2007 | 0.50 | 1.45 | 7.10 | 7.47 |
| 1968 | 1.17 | 7.66 | 3.38 | NA | 2008 | 0.45 | 1.04 | 7.46 | 9.90 |
| 1969 | 1.58 | 7.38 | 3.38 | NA | 2009 | 0.35 | 1.22 | 7.87 | 8.84 |
| 1970 | 1.37 | 7.19 | 3.55 | NA | 2010 | 0.30 | 1.41 | 7.39 | 3.63 |
| 1971 | 1.17 | 7.83 | 3.73 | NA | 2011 | 0.27 | 1.31 | 7.39 | 6.62 |
| 1972 | 1.38 | 7.09 | 3.90 | NA | 2012 | 0.27 | 1.31 | 7.39 | NA |
| 1973 | 0.99 | 6.63 | 4.09 | NA | | | | | |
| 1974 | 0.72 | 7.01 | 4.26 | NA | | | | | |

APPENDIX B ESTIMATION OF ABUNDANCE INDICES FROM THE IPHC SURVEY

Annual indices of catch rate (CPUE) in any year y were obtained by taking the overall mean of the mean Bocaccio CPUE in each of the surveyed strata i :

$$U_y = \frac{\sum_{i=1}^{k_y} C_{yi}}{k_y} \quad \text{Eq. 1}$$

where C_{yi} = mean CPUE (pieces/skate) for Bocaccio in year y in stratum i ;

k_y = number of strata in year y ;

U_y = mean CPUE of Bocaccio for year y .

CPUE (C_{yi}) in stratum i for year y was calculated as pieces per skate by

$$C_{yi} = \frac{\sum_{j=1}^{n_{yi}} \left(\frac{P_{yij}}{S_{yij}} \right)}{n_{yi}} \quad \text{Eq. 2}$$

where P_{yij} = number of pieces of Bocaccio in year y in stratum i and set j ;

S_{yij} = number of skates in year y by set j in stratum i ;

n_{yi} = number of sets in year y for stratum i .

CPUE estimates were bootstrapped for 1000 random draws with replacement to obtain bias- corrected (Efron 1982) 95% confidence regions for each year.

APPENDIX C RELATIVE ABUNDANCE INDICES

Appendix Table 4. Arithmetic and standardised commercial bottom trawl CPUE indices with upper and lower bounds of the standardised indices and the associated standard error for the 3C-5E model of non-zero catches of Bocaccio. The geometric mean of the standardised series has been scaled so that it equals the geometric mean of the arithmetic series

| Fishing year | Arithmetic | Standardised | Lower bound | Upper bound | Standard error |
|---------------------|-------------------|---------------------|--------------------|--------------------|-----------------------|
| 96/97 | 29.8 | 28.9 | 27.1 | 30.8 | 0.032 |
| 97/98 | 29.4 | 31.5 | 30.0 | 33.1 | 0.025 |
| 98/99 | 27.4 | 27.9 | 26.6 | 29.3 | 0.025 |
| 99/00 | 25.2 | 27.4 | 26.2 | 28.7 | 0.024 |
| 00/01 | 32.1 | 28.1 | 26.9 | 29.3 | 0.022 |
| 01/02 | 33.5 | 32.3 | 30.9 | 33.8 | 0.022 |
| 02/03 | 29.4 | 29.9 | 28.6 | 31.2 | 0.022 |
| 03/04 | 27.1 | 27.9 | 26.7 | 29.2 | 0.023 |
| 04/05 | 26.0 | 21.9 | 20.9 | 23.0 | 0.025 |
| 05/06 | 18.9 | 20.5 | 19.5 | 21.5 | 0.024 |
| 06/07 | 18.2 | 19.5 | 18.4 | 20.6 | 0.028 |

Appendix Table 5. Biomass estimates for Bocaccio from the QCSd Shrimp Trawl Survey for the survey years 1999 to 2011. Bootstrap bias corrected confidence intervals and CVs are based on 1,000 random draws with replacement. The analytic CV is based on the assumption of random tow selection within a stratum. – indicates not applicable.

| Survey Year | Biomass (t) | Mean bootstrap biomass (t) | Lower bound biomass (t) | Upper bound biomass (t) | Bootstrap CV | Analytic CV |
|--------------------|--------------------|-----------------------------------|--------------------------------|--------------------------------|---------------------|--------------------|
| 1999 | 18.8 | 19.0 | 5.3 | 38.4 | 0.432 | 0.445 |
| 2000 | 9.2 | 9.3 | 0.0 | 29.1 | 0.796 | 0.761 |
| 2001 | 19.4 | 19.5 | 5.7 | 39.7 | 0.432 | 0.420 |
| 2002 | 2.5 | 2.6 | 0.0 | 10.3 | 0.980 | 1.000 |
| 2003 | 7.2 | 7.5 | 0.0 | 17.0 | 0.557 | 0.571 |
| 2004 | 17.7 | 17.5 | 0.0 | 51.8 | 0.840 | 0.865 |
| 2005 | 4.7 | 4.4 | 0.0 | 19.1 | 1.014 | 1.000 |
| 2006 | 7.1 | 7.0 | 1.6 | 16.2 | 0.522 | 0.532 |
| 2007 | 0.0 | 0.0 | – | – | – | 0.000 |
| 2008 | 0.0 | 0.0 | – | – | – | 0.000 |
| 2009 | 10.9 | 10.8 | 3.6 | 21.1 | 0.417 | 0.413 |
| 2010 | 0.0 | 0.0 | – | – | – | 0.000 |
| 2011 | 462.6 | 467.8 | 0.0 | 1,946.0 | 0.988 | 1.000 |

Appendix Table 6. Biomass estimates for Bocaccio from the WCVI shrimp trawl survey for the survey years 1975 to 2011. Biomass estimates are based on a post-stratification of this survey into two strata and by assuming that the survey tows were randomly selected within these areas. Bootstrap bias corrected confidence intervals and CVs are based on 1,000 random draws with replacement. The analytic CV is based on the assumption of random tow selection within a stratum. – indicates not applicable

| Survey Year | Biomass (t) | Mean bootstrap biomass (t) | Lower bound biomass (t) | Upper bound biomass (t) | Bootstrap CV | Analytic CV |
|-------------|-------------|----------------------------|-------------------------|-------------------------|--------------|-------------|
| 1975 | 106.1 | 107.0 | 48.7 | 190.9 | 0.340 | 0.350 |
| 1976 | 42.3 | 42.3 | 11.5 | 99.4 | 0.508 | 0.521 |
| 1977 | 84.7 | 84.6 | 28.4 | 177.1 | 0.449 | 0.467 |
| 1978 | 362.1 | 357.3 | 8.5 | 1,000.2 | 0.715 | 0.713 |
| 1979 | 25.6 | 25.6 | 5.1 | 52.9 | 0.456 | 0.494 |
| 1980 | 21.2 | 20.8 | 0.0 | 58.2 | 0.735 | 0.768 |
| 1981 | 28.6 | 28.6 | 0.7 | 89.5 | 0.752 | 0.781 |
| 1982 | 577.0 | 581.6 | 54.0 | 1,741.1 | 0.821 | 0.823 |
| 1983 | 339.6 | 352.4 | 7.3 | 1,293.4 | 0.920 | 0.926 |
| 1985 | 366.9 | 368.2 | 168.6 | 606.0 | 0.301 | 0.302 |
| 1987 | 73.7 | 73.5 | 26.6 | 138.9 | 0.379 | 0.380 |
| 1988 | 117.9 | 115.0 | 25.7 | 275.7 | 0.537 | 0.525 |
| 1989 | 33.6 | 33.3 | 7.0 | 89.8 | 0.558 | 0.531 |
| 1990 | 162.6 | 163.5 | 30.0 | 421.3 | 0.612 | 0.591 |
| 1991 | 115.3 | 115.3 | 5.4 | 395.0 | 0.826 | 0.903 |
| 1992 | 387.0 | 379.6 | 111.6 | 854.0 | 0.449 | 0.426 |
| 1993 | 10.0 | 10.1 | 0.0 | 40.9 | 1.001 | 1.000 |
| 1994 | 139.6 | 138.5 | 0.0 | 535.3 | 0.958 | 0.945 |
| 1995 | 15.4 | 15.1 | 0.0 | 59.2 | 0.991 | 1.000 |
| 1996 | 50.5 | 50.2 | 0.0 | 174.2 | 0.870 | 0.902 |
| 1997 | 110.9 | 111.0 | 21.4 | 267.0 | 0.575 | 0.576 |
| 1998 | 214.3 | 212.2 | 0.0 | 729.4 | 0.909 | 0.940 |
| 1999 | 2.0 | 2.0 | 0.0 | 7.0 | 0.951 | 1.000 |
| 2000 | 0.0 | 0.0 | – | – | – | 0.000 |
| 2001 | 70.2 | 69.5 | 19.4 | 156.3 | 0.468 | 0.460 |
| 2002 | 30.6 | 30.7 | 1.0 | 93.5 | 0.758 | 0.765 |
| 2003 | 32.1 | 32.3 | 0.0 | 72.5 | 0.530 | 0.552 |
| 2004 | 30.2 | 29.7 | 0.0 | 88.9 | 0.731 | 0.726 |
| 2005 | 583.2 | 570.8 | 0.0 | 2,050.1 | 0.976 | 0.971 |
| 2006 | 6.4 | 6.5 | 0.0 | 26.8 | 0.977 | 1.000 |
| 2007 | 11.6 | 11.3 | 0.3 | 37.5 | 0.732 | 0.693 |
| 2008 | 16.1 | 16.0 | 0.0 | 36.6 | 0.569 | 0.586 |
| 2009 | 91.1 | 92.5 | 19.7 | 181.4 | 0.452 | 0.461 |
| 2010 | 47.3 | 46.6 | 8.4 | 112.1 | 0.561 | 0.563 |
| 2011 | 0.0 | 0.0 | – | – | – | 0.000 |

Appendix Table 7. Biomass estimates for Bocaccio from the West Coast Haida Gwaii groundfish synoptic trawl survey for the years 2006 to 2010. Biomass estimates are based on a post-stratification of this survey into two strata and by assuming that the survey tows were randomly selected within these areas. Bootstrap bias corrected confidence intervals and CVs are based on 1,000 random draws with replacement. The analytic CV is based on the assumption of random tow selection within a stratum.

| Survey Year | Biomass (t) | Mean bootstrap biomass | Lower bound biomass (t) | Upper bound biomass (t) | Bootstrap CV | Analytic CV |
|-------------|-------------|------------------------|-------------------------|-------------------------|--------------|-------------|
| 2006 | 9.9 | 10.0 | 4.3 | 17.1 | 0.329 | 0.345 |
| 2007 | 9.6 | 9.6 | 4.3 | 16.9 | 0.328 | 0.329 |
| 2008 | 12.0 | 12.0 | 6.0 | 20.4 | 0.309 | 0.301 |
| 2010 | 8.0 | 8.2 | 3.4 | 14.5 | 0.352 | 0.359 |

Appendix Table 8. Biomass estimates for Bocaccio from the Hecate Strait Groundfish synoptic trawl survey for the years 2005 to 2011. Biomass estimates are based on a post-stratification of this survey into two strata and by assuming that the survey tows were randomly selected within these areas. Bootstrap bias corrected confidence intervals and CVs are based on 1,000 random draws with replacement. The analytic CV is based on the assumption of random tow selection within a stratum.

| Survey Year | Biomass (t) | Mean bootstrap biomass (t) | Lower bound biomass (t) | Upper bound biomass (t) | Bootstrap CV | Analytic CV |
|-------------|-------------|----------------------------|-------------------------|-------------------------|--------------|-------------|
| 2005 | 19.5 | 19.4 | 8.3 | 36.4 | 0.376 | 0.369 |
| 2007 | 48.6 | 48.7 | 15.6 | 95.7 | 0.403 | 0.389 |
| 2009 | 16.8 | 16.7 | 5.5 | 35.7 | 0.450 | 0.445 |
| 2011 | 55.1 | 55.3 | 6.8 | 152.1 | 0.633 | 0.621 |

Appendix Table 9. Biomass estimates for Bocaccio from the Queen Charlotte Sound Groundfish synoptic trawl survey for the years 2005 to 2011. Biomass estimates are based on a post-stratification of this survey into two strata and by assuming that the survey tows were randomly selected within these areas. Bootstrap bias corrected confidence intervals and CVs are based on 1,000 random draws with replacement. The analytic CV is based on the assumption of random tow selection within a stratum.

| Survey Year | Biomass (t) | Mean bootstrap biomass (t) | Lower bound biomass (t) | Upper bound biomass (t) | Bootstrap CV | Analytic CV |
|-------------|-------------|----------------------------|-------------------------|-------------------------|--------------|-------------|
| 2003 | 110.1 | 109.5 | 26.4 | 271.0 | 0.591 | 0.606 |
| 2004 | 308.9 | 303.6 | 46.5 | 912.2 | 0.788 | 0.776 |
| 2005 | 295.0 | 302.9 | 57.8 | 849.7 | 0.692 | 0.704 |
| 2007 | 127.8 | 126.3 | 28.7 | 351.1 | 0.640 | 0.647 |
| 2009 | 88.5 | 92.9 | 20.1 | 218.0 | 0.585 | 0.613 |
| 2011 | 36.0 | 36.6 | 12.7 | 75.6 | 0.439 | 0.436 |

Appendix Table 10. Biomass estimates for Bocaccio from the West Coast Vancouver Island Groundfish synoptic trawl survey for the years 2006 to 2010. Biomass estimates are based on a post-stratification of this survey into two strata and by assuming that the survey tows were randomly selected within these areas. Bootstrap bias corrected confidence intervals and CVs are based on 1,000 random draws with replacement. The analytic CV is based on the assumption of random tow selection within a stratum

| Survey Year | Biomass (t) | Mean bootstrap biomass (t) | Lower bound biomass (t) | Upper bound biomass (t) | Bootstrap CV | Analytic CV |
|-------------|-------------|----------------------------|-------------------------|-------------------------|--------------|-------------|
| 2004 | 370.8 | 390.0 | 40.4 | 1149.2 | 0.760 | 0.783 |
| 2006 | 336.1 | 337.1 | 69.2 | 989.0 | 0.715 | 0.705 |
| 2008 | 155.1 | 155.9 | 88.3 | 255.4 | 0.270 | 0.278 |
| 2010 | 53.2 | 53.6 | 22.1 | 97.7 | 0.371 | 0.385 |

Appendix Table 11. Biomass estimates for Bocaccio in the U.S. Triennial survey (Canadian waters only) with 95% confidence regions based on the bootstrap distribution of biomass. Biomass estimates are calculated as described earlier. The bootstrap estimates are based on 5000 random draws with replacement (from Stanley et al. 2009a).

| Estimate type | Year | Biomass | Mean bootstrap biomass | Lower bound biomass | Upper bound biomass | CV bootstrap | CV Analytic |
|---------------|------|---------|------------------------|---------------------|---------------------|--------------|-------------|
| Canada | 1980 | 8,103 | 8,261 | 296 | 30,812 | 0.923 | 0.937 |
| Vancouver | 1983 | 4,731 | 4,611 | 681 | 14,566 | 0.697 | 0.688 |
| | 1989 | 1,279 | 1,302 | 338 | 2,657 | 0.454 | 0.456 |
| | 1992 | 792 | 797 | 135 | 2,149 | 0.633 | 0.654 |
| | 1995 | 65 | 64 | 16 | 135 | 0.448 | 0.467 |
| | 1998 | 141 | 140 | 49 | 279 | 0.409 | 0.408 |
| | 2001 | 120 | 123 | 0 | 365 | 0.768 | 0.798 |


Appendix Table 12. Estimates and 95% confidence limits of relative catch rate (pieces/skate) of Bocaccio in the IPHC BC longline survey

| Survey Year | CPUE | | |
|-------------|----------------|-------------|-------------|
| | Bootstrap Mean | Lower bound | Upper bound |
| 2003 | 0.013 | 0.006 | 0.024 |
| 2004 | 0.023 | 0.009 | 0.038 |
| 2005 | 0.013 | 0.005 | 0.024 |
| 2006 | 0.036 | 0.010 | 0.079 |
| 2007 | 0.018 | 0.008 | 0.028 |
| 2008 | 0.038 | 0.019 | 0.062 |
| 2009 | 0.020 | 0.009 | 0.034 |
| 2010 | 0.011 | 0.004 | 0.021 |
| 2011 | 0.022 | 0.008 | 0.039 |

APPENDIX D ADDITIONAL REFERENCE CASE RESULTS

Appendix Table 13. Posterior means, medians, standard deviations (SD), CVs and 95% probability intervals for q-gross (qgfin). The last three columns show the 2.5th, 50th, and 97.5th percentiles of the random variable qgfin. The mean and SD of the natural logarithm of qgfin were used as inputs to the multivariate log normal prior density function for the survey q parameter in the stock assessment.

| Survey | Mean | SD | CV | 2.5% | Median | 97.5% |
|------------------------------|-------------|-----------|-----------|-------------|---------------|--------------|
| #1 - WCVI groundfish | 0.11100 | 0.0760 | 0.69 | 0.017000 | 0.090000 | 0.3050 |
| #2 - QCSd-groundfish | 0.07200 | 0.0520 | 0.72 | 0.010500 | 0.058000 | 0.2060 |
| #3 - HS – groundfish | 0.01050 | 0.0080 | 0.76 | 0.001500 | 0.008300 | 0.0315 |
| #4 - WCHG - groundfish | 0.00340 | 0.0024 | 0.71 | 0.000490 | 0.002700 | 0.0096 |
| #5 - WCVI Shrimp | 0.00480 | 0.0066 | 1.40 | 0.000300 | 0.002600 | 0.0222 |
| #6 - QCSd Shrimp | 0.00046 | 0.0012 | 2.64 | 0.000001 | 0.000107 | 0.0032 |
| #7 - US Triennial groundfish | 0.06050 | 0.0945 | 1.56 | 0.000600 | 0.024000 | 0.3360 |



Appendix Figure 1. Marginal density functions for q -gross ($qgfin$) for the seven different surveys when Bayesian updating and uncertainty factors are applied to the q -net factors